

RESEARCH

Open Access

# NO<sub>2</sub> inhalation induces maturation of pulmonary CD11c<sup>+</sup> cells that promote antigen-specific CD4<sup>+</sup> T cell polarization

Samantha R Hodgkins<sup>1</sup>, Jennifer L Ather<sup>1</sup>, Sara A Paveglio<sup>1</sup>, Jenna L Allard<sup>1</sup>, Laurie A Whittaker LeClair<sup>1</sup>, Benjamin T Suratt<sup>1</sup>, Jonathan E Boyson<sup>2</sup>, Matthew E Poynter<sup>1\*</sup>

## Abstract

**Background:** Nitrogen dioxide (NO<sub>2</sub>) is an air pollutant associated with poor respiratory health, asthma exacerbation, and an increased likelihood of inhalational allergies. NO<sub>2</sub> is also produced endogenously in the lung during acute inflammatory responses. NO<sub>2</sub> can function as an adjuvant, allowing for allergic sensitization to an innocuous inhaled antigen and the generation of an antigen-specific Th2 immune response manifesting in an allergic asthma phenotype. As CD11c<sup>+</sup> antigen presenting cells are considered critical for naïve T cell activation, we investigated the role of CD11c<sup>+</sup> cells in NO<sub>2</sub>-promoted allergic sensitization.

**Methods:** We systemically depleted CD11c<sup>+</sup> cells from transgenic mice expressing a simian diphtheria toxin (DT) receptor under of control of the CD11c promoter by administration of DT. Mice were then exposed to 15 ppm NO<sub>2</sub> followed by aerosolized ovalbumin to promote allergic sensitization to ovalbumin and were studied after subsequent inhaled ovalbumin challenges for manifestation of allergic airway disease. In addition, pulmonary CD11c<sup>+</sup> cells from wildtype mice were studied after exposure to NO<sub>2</sub> and ovalbumin for cellular phenotype by flow cytometry and *in vitro* cytokine production.

**Results:** Transient depletion of CD11c<sup>+</sup> cells during sensitization attenuated airway eosinophilia during allergen challenge and reduced Th2 and Th17 cytokine production. Lung CD11c<sup>+</sup> cells from wildtype mice exhibited a significant increase in MHCII, CD40, and OX40L expression 2 hours following NO<sub>2</sub> exposure. By 48 hours, CD11c<sup>+</sup>MHCII<sup>+</sup> DCs within the mediastinal lymph node (MLN) expressed maturation markers, including CD80, CD86, and OX40L. CD11c<sup>+</sup>CD11b<sup>-</sup> and CD11c<sup>+</sup>CD11b<sup>+</sup> pulmonary cells exposed to NO<sub>2</sub> *in vivo* increased uptake of antigen 2 hours post exposure, with increased ova-Alexa 647<sup>+</sup> CD11c<sup>+</sup>MHCII<sup>+</sup> DCs present in MLN from NO<sub>2</sub>-exposed mice by 48 hours. Co-cultures of ova-specific CD4<sup>+</sup> T cells from naïve mice and CD11c<sup>+</sup> pulmonary cells from NO<sub>2</sub>-exposed mice produced IL-1, IL-12p70, and IL-6 *in vitro* and augmented antigen-induced IL-5 production.

**Conclusions:** CD11c<sup>+</sup> cells are critical for NO<sub>2</sub>-promoted allergic sensitization. NO<sub>2</sub> exposure causes pulmonary CD11c<sup>+</sup> cells to acquire a phenotype capable of increased antigen uptake, migration to the draining lymph node, expression of MHCII and co-stimulatory molecules required to activate naïve T cells, and secretion of polarizing cytokines to shape a Th2/Th17 response.

## Background

The prevalence of allergic asthma has risen steadily in recent decades, making the disease a primary public health concern [1]. Potential explanations for the increase include reduced exposure to infectious agents

during childhood, dietary changes, and exposure to environmental pollutants. Allergic asthma is caused primarily by an inappropriate CD4<sup>+</sup> Th2 response, which results in symptoms mediated by Th2 cytokines, including IL-13 provoking airways hyperresponsiveness and mucus production, IL-4 promoting the production of antigen specific IgE, and IL-5 inducing eosinophilia [2]. Recent evidence suggests that Th17 cells secreting IL-17

\* Correspondence: [matthew.poynter@uvm.edu](mailto:matthew.poynter@uvm.edu)

<sup>1</sup>The Vermont Lung Center and Department of Medicine, University of Vermont, Burlington, VT, 05405, USA

are associated with a severe [3], steroid-resistant [4] form of allergic asthma. However, the underlying mechanisms that initiate the aberrant T cell response in allergic asthma are still not well understood (reviewed in [5]). Our lab has shown that inhalation of the gaseous air pollutant and endogenously-generated reactant nitrogen dioxide (NO<sub>2</sub>) is capable of acting as an adjuvant, promoting allergic sensitization to the innocuous protein ovalbumin (ova) in a novel mouse model [6]. This model is physiologically relevant as antigen sensitization occurs via inhalation, as would typically occur in humans and does not require an additional adjuvant [7]. NO<sub>2</sub> has also been correlated with poor respiratory health [8], exacerbating existing asthma in animal models [9] and in human subjects [10], as well as with an increased likelihood of inhalational allergies [11] and developing asthma in human studies [12].

Pulmonary antigen-presenting cells, especially dendritic cells (DCs), express the surface marker CD11c [13] and have a potent ability to induce the proliferation and activation of naïve T cells and to secrete inflammatory and T-helper cell polarizing cytokines [14-16]. CD11c<sup>+</sup> cells are critical for initiating and shaping the antigen-specific adaptive immune response and are critical during the reactivation of CD4<sup>+</sup> T cells *in vivo* [17]. CD11c<sup>+</sup> DCs are capable of these activities because they possess multiple unique characteristics. First, DCs are strategically located beneath the airway epithelium and continually take up antigen under steady-state conditions [15]. Second, DCs can undergo maturation upon exposure to inflammatory stimuli and travel to draining lymph nodes, presenting antigens in the context of both MHC I and MHC II. Finally, DCs express co-stimulatory molecules and secrete polarizing cytokines necessary to initiate and shape the T cell mediated immune response [16,18]. However, defining DCs via surface marker expression remains complicated, especially in non-lymphoid tissues such as the lung, due to the number of different methods described in the literature and the shared cell surface markers expressed by several cell subsets. The myeloid DC subset is attributed with T cell stimulatory capacity, having the ability to induce Th1, Th2, or Th17 type responses [19], as well as non-inflammatory T regulatory (Treg) responses [20]. Myeloid DCs in the lung have been defined as CD11c<sup>+</sup>CD11b<sup>+</sup> [19,21], CD11c<sup>+</sup>MHCII<sup>+</sup> [22], or CD11c<sup>+</sup> alone or in combination with low FITC auto-fluorescence [23,24]. This variation is further complicated by the overlap of markers with multiple other cell types, the most prominent of which in the lung is CD11c<sup>+</sup> macrophages [21]. Plasmacytoid DCs (pDCs) are also present within the lung [25] and have been shown to exert an anti-inflammatory role, decreasing both the ability of mDCs to generate effector T cells as well as inducing Treg cell

proliferation [26]. This pDC population expresses B220 (CD45RB) as well as low levels of Gr-1, making them more difficult to delineate from B cells and granulocytes than myeloid DCs [18]. Recently, another population of dendritic cells called inflammatory DCs has been described (CD11b<sup>+</sup>Gr-1<sup>lo</sup>F4/80<sup>lo</sup>), which traffic from the blood to sites of inflammation, upregulate CD11c upon arrival in the tissue, and acquire the DC characteristics of migration to the draining lymph node and induction of T cell proliferation [22,27].

DCs are capable of skewing the T helper cell response through their ability to express distinct patterns of co-stimulatory molecules as well as to produce cytokines that create an environment for differential T cell polarization [19]. Expression of the co-stimulatory molecules CD86 and OX40L have been shown to promote naïve CD4<sup>+</sup> T cells to develop a Th2 phenotype [19,28,29]. Importantly, OX40L-deficient mice are protected from allergic sensitization and Th2 mediated inflammation, indicating that OX40L plays a critical role in the generation of Th2 immune responses [29]. DCs also regulate Th2 cell differentiation and expansion by producing IL-6 [30,31]. Th17 cells may also be induced by DC production of IL-6 in combination with TGFβ or IL-23 [30,32,33], or by IL-1β alone [34], while IL-12 alone promotes a Th1 response [35]. Thus, activated DCs as a cellular source of co-stimulatory molecules and polarizing cytokines function as important regulators of CD4<sup>+</sup> mediated T cell responses.

Since activation of pathogenic CD4<sup>+</sup> Th2 cells is believed to be dependent on CD11c<sup>+</sup> DCs [36], we tested whether CD11c<sup>+</sup> cells are necessary for NO<sub>2</sub>-promoted allergic sensitization. We used transgenic mice in which CD11c<sup>+</sup> cells can be temporarily depleted due to the incorporation of a transgene encoding a simian Diphtheria Toxin Receptor (DTR) and Green Fluorescent Protein (GFP) fusion protein under the control of the murine CD11c promoter (CD11c-DTR mice) [37,38]. As murine cells lack the DT receptor [17] and only CD11c<sup>+</sup> cells in CD11c-DTR Tg<sup>+</sup> mice express the DT receptor, administration of DT to these mice results in inducible ablation of 90% of CD11c<sup>+</sup> cells within the mouse lasting for approximately 48 hours [38]. Using CD11c-DTR mice, it has been demonstrated in models of allergic asthma that CD11c<sup>+</sup> DCs are important during allergic sensitization induced by the i.p. injection of ova in combination with the adjuvant Alum [22] and are also critical during allergen challenge [17]. Importantly, the use of these mice to assess the role of CD11c<sup>+</sup> cells during allergic sensitization via inhalation, and in response to NO<sub>2</sub>, is so far unpublished.

In these studies, we demonstrate that when CD11c<sup>+</sup> cells were depleted during sensitization, mice exhibited less inflammation within the lung following allergen challenge

and displayed a significantly impaired capacity for CD4<sup>+</sup> T cells to secrete Th2 and Th17 cytokines *in vitro*, implying that CD11c<sup>+</sup> cells are critical during NO<sub>2</sub>-promoted allergic sensitization. Furthermore, flow cytometric analyses of CD11c<sup>+</sup> cell populations within the lung and draining lymph node revealed that NO<sub>2</sub> increased MHCII and costimulatory molecule expression in a temporally-orchestrated fashion. Finally, CD11c<sup>+</sup> pulmonary cells exposed to NO<sub>2</sub> increased antigen capture of ova-Alexa 647, produced significant amounts of IL-1 $\alpha$ , IL-1 $\beta$ , IL-12p70, and IL-6 compared to air controls, and augmented the production of IL-5 by CD4<sup>+</sup> T cells *in vitro*. These results suggest a critical function for pulmonary CD11c<sup>+</sup> cells during NO<sub>2</sub>-promoted allergic sensitization, including augmented pulmonary CD11c<sup>+</sup> cell recruitment and an improved ability to uptake and present antigen, provide co-stimulation, and secrete polarizing cytokines that initiate and shape a Th2/Th17-biased immune response. Comprehending the mechanisms underlying NO<sub>2</sub>-promoted allergic sensitization provides insight into the potential for common environmental pollutants to affect the development of increasingly prevalent pulmonary disorders, such as allergic asthma.

## Methods

### Mice

C57BL/6 mice were purchased from the Jackson Laboratory and used in experiments and as antigen presenting cell donors. OTII TCR transgenic mice on the C57BL/6 background were purchased from the Jackson Laboratory and bred at the University of Vermont as a source of CD4 T cells responsive to the peptide ova<sub>323-339</sub>, an immunodominant MHCII antigenic epitope from the protein ovalbumin [39]. CD11c-DTR-EGFP (CD11c-DTR) transgenic mice were purchased from Jackson Laboratories (Bar Harbor, ME) (B6.FVB-Tg(Itgax-DTR/EGFP)57Lan/J). The CD11c-DTR mice were generated with a transgene insert containing the Itgax (CD11c) promoter driving expression of a fusion protein containing the diphtheria toxin receptor (DTR) and the enhanced green fluorescent protein (GFP). The CD11c-DTR transgenic and transgene-negative littermates used in these studies were bred and raised at the University of Vermont. Eight- to fifteen-week old female mice, weighing approximately 20-25 g, were used for all studies. Mice were euthanized by a lethal dose (150 mg/kg) of pentobarbital (Nembutal, Ovation Pharmaceuticals Inc., Deerfield, IL) via intraperitoneal (i.p.) injection. All experiments were conducted following the guidelines and under the approval of the University of Vermont's Institutional Animal Care and Use Committee. The mice were housed in a facility accredited by the American Association for the Accreditation of Laboratory Animal Care.

### In vivo NO<sub>2</sub> and ovalbumin exposures

1000 ppm nitrogen dioxide with a nitrogen balance (Air-Gas, Salem, NH) was diluted to 15 ppm with HEPA-filtered room air in a specially designed glass chamber (Specialty Glass, Inc., Rosharon, TX) within a fume hood. As needed, control mice were exposed to HEPA-filtered room air in an identical chamber for the same duration as NO<sub>2</sub>-exposed mice. A calibrated nitric oxide analyzer, equipped with an NO<sub>2</sub> thermal converter, was used to measure NO<sub>2</sub> in the gas phase, according to manufacturer's instructions (EcoPhysics, Ann Arbor, MI). For studies in which exposure to ovalbumin was performed, mice were exposed to aerosolized 3.4% ovalbumin (final concentration 10 mg/ml) (Grade V, Sigma, St. Louis, MO) in Dulbecco's Phosphate-Buffered Saline (DPBS) (CellGro, Manassas, VA) for 30 minutes immediately following exposure to either NO<sub>2</sub> or air or during antigen challenge. The contaminating endotoxin concentration in the ovalbumin preparation used for inhalation was measured to be 1.5 ng/mg protein using a chromogenic Limulus assay (GenScript, Piscataway, NJ).

### NO<sub>2</sub>-promoted allergic sensitization model

Eight-week-old C57BL/6 female mice or eight to fifteen-week-old male and female CD11c-DTR mice (Tg<sup>+</sup> and Tg<sup>-</sup>) were exposed to 15 ppm NO<sub>2</sub> for 1 hour followed by 30 minutes of aerosolized ova on day 0. This acute pro-inflammatory concentration of NO<sub>2</sub> was established in a separate dose response study to be the minimum sufficient to promote allergic sensitization using this regimen. Mice were then challenged with aerosolized ova on days 14, 15, and 16 and analyzed 48 hours later (day 18). As previously described [6], mice exposed to NO<sub>2</sub> and saline during the sensitization phase do not elicit allergic responses and were only included in limited numbers as controls to assure that in wildtype mice the NO<sub>2</sub>/ova exposure regimen induced airway eosinophilia, elevated levels of antigen-specific immunoglobulins in the serum, and antigen-specific cytokine production during restimulation of CD4<sup>+</sup> T cells *in vitro* compared to these controls.

### Administration of diphtheria toxin

As dictated by the experiment, Tg<sup>+</sup> and Tg<sup>-</sup> CD11c-DTR mice were administered Diphtheria toxin (DT) (Sigma, St. Louis, MO) or saline via i.p. injection 24 hours prior to either sensitization or flow cytometric analyses for depletion of CD11c<sup>+</sup> cells. A dose of 4 ng/g of body weight was used at a concentration of 0.2 or 0.25 ng/ $\mu$ l.

### Bronchoalveolar lavage (BAL)

Following euthanasia, BAL was collected by instilling and recovering 1 ml of DPBS (CellGro, Manassas, VA)

containing protease inhibitor cocktail (Sigma, St. Louis, MO) into the lung via a tracheal cannula. Total and differential cell counts were performed using the Advia 120 automated hematology analyzer system and cytopins. For the cytopins, cells were centrifuged onto glass slides at 800 rpm (RCF =  $80 \times g$ ) and stained using the Hema3 kit (Biochemical Sciences, Inc., Swedesboro, NJ), with differential cell counts performed on at least 200 cells [40]. BALF supernatants were collected for protein quantitation by Bradford assay (Bio-Rad, Hercules, CA) and cytokine quantitation by Bio-Plex (Bio-Rad).

#### **Ova-specific IgE and IgG<sub>1</sub> quantification**

Following euthanasia, approximately 300  $\mu$ l of blood was collected via cardiac puncture of the right ventricle using a 26-gauge needle attached to a 1 ml syringe into serum separator tubes (BD Biosciences, San Diego, CA). The blood was centrifuged at 13,200 rpm and serum was stored at  $-80^{\circ}\text{C}$ . Ova-specific immunoglobulins were quantified utilizing a two-step sandwich (capture) ELISA. 96-well high-binding plates (Costar, Bethesda, MD) were coated with 2  $\mu$ g/ml anti-mouse IgE (BD Pharmingen clone R35-72) or IgG<sub>1</sub> (BD Pharmingen clone A85-3) mAb in DPBS for 1 hour at RT. Plates were washed with 0.05% Tween 20 (Fisher Scientific, Pittsburgh, PA) in PBS, blocked for 1 hour with PBS/1% BSA (Fischer Scientific), and serum samples were diluted (1:10 for IgE, 1:1000 for IgG<sub>1</sub>) and added in duplicate in PBS/1% BSA for 1 hour at RT. Plates were washed and incubated with a 1:2500 dilution of digoxigenin-coupled ova (Roche, Madison, WI) in PBS/1% BSA for 1 hour at RT. Plates were washed and incubated with a 1:2000 dilution of anti-digoxigenin-Fab coupled to peroxidase (Roche) in PBS/1% BSA for 30 minutes. Plates were washed, developed using reagents from R&D Systems (Minneapolis, MN), stopped with 1 N H<sub>2</sub>SO<sub>4</sub>, and ODs were read using a Bio-Tek Instruments PowerWave<sub>x</sub> at 450 nm with background subtraction at 570 nm.

#### **CD4<sup>+</sup> T cell re-stimulation and cytokine analyses**

Single-cell suspensions were generated from spleens and mediastinal lymph nodes (MLNs) by passing the tissues through a 70  $\mu$ m nylon mesh filter (BD Biosciences) and lymphocytes were enriched by separation with Lymphocyte Separation Medium (MP Biomedicals, Irvine, CA). CD4<sup>+</sup> T cells were isolated by positive selection using CD4 magnetic microbeads (Miltenyi Biotec, Bergisch Gladbach, Germany), according to the manufacturer's instructions. CD4<sup>+</sup> T cells ( $2 \times 10^6$  cells/ml) were stimulated with 100  $\mu$ g/ml ova in the presence of C57BL/6 antigen presenting cells (APCs) ( $4 \times 10^6$  cells/ml). APCs were obtained by splenic T cell depletion by negative selection using Abs to CD4, CD8, and Thy-1, and

treatment with rabbit complement and mitomycin C, as previously described [41]. Following 96 hours of stimulation, supernatants were collected and analyzed by ELISA using reagents and instructions from R&D Systems. ODs from duplicate samples and duplicate standards were read using a Bio-Tek Instruments PowerWave<sub>x</sub> at 450 nm with background subtraction at 570 nm.

#### **Analysis of CD11c<sup>+</sup> cells from the lung and mediastinal lymph node (MLN)**

The lung was dissociated to a single cell suspension by mechanical disruption followed by incubation at  $37^{\circ}\text{C}$  with 0.5 mg/ml Liberase Blendzyme 3 (Roche Applied Science, Indianapolis, IN) and 40  $\mu$ g/ml DNase for a total of 40 minutes. Red blood cells were lysed using 0.83% ammonium chloride and the lung suspension was filtered through a 40  $\mu$ m nylon mesh membrane (BD Biosciences, San Diego, CA). Total cells were then counted using Advia or hemocytometer and were stained, as described below, or used for the isolation of CD11c<sup>+</sup> pulmonary DCs via positive selection using MACS columns (Miltenyi Biotec, Bergisch Gladbach, Germany) for *in vitro* studies, according to the manufacturer's instructions. MLN cells were dissociated by mechanical disruption, filtered through a 40  $\mu$ m nylon mesh membrane (BD Biosciences), and stained as described below.

#### **Cell staining and fluorescence-activated cell scanning (FACScan)**

Lung and MLN derived cells were stained with the following antibodies: CD45-PO, CD11c-PETR, F4/80-Alexa 647, CD86-Alexa 647 (all from Caltag, Carlsbad, CA); I-A/I-E-PerCP/Cy5.5 (BioLegend, San Diego, CA); CD11b-APCcy7 and GR-1-PE (all from BD Pharmingen, San Diego, CA). To assess maturation, cells were also stained for CD80-PE, CD40-APC, and biotinylated OX40L (all from BD Pharmingen). Biotinylated antibodies were detected using streptavidin-PE (BD Pharmingen).  $1 \times 10^6$  cells were first blocked with 2.5  $\mu$ g/ml Fc block (anti-CD16/CD32, BD Pharmingen) for 30 minutes, washed in FACS buffer (DPBS (CellGro) with 5% FBS (Gibco, Carlsbad, CA)), and then stained for 30 minutes in 100  $\mu$ l of antibody solution at the optimal concentration. In the case of a biotinylated primary antibody, cells were washed and stained with the secondary antibody for 30 minutes. Following staining, all cells were washed and fixed in DPBS with 5% FBS and 1% paraformaldehyde. Cells were analyzed on the flow cytometer 1-3 days following staining using a Becton Dickinson LSR II FACS equipped to distinguish as many as 7 fluorophores. Dead cells were excluded from analysis by FSC and SSC gating.

### Oropharyngeal aspiration of ova-Alexa 647

Following exposure to 1 hour of air or 15 ppm NO<sub>2</sub>, mice were anesthetized with inhaled isoflurane (Webster Veterinary, Sterling, MA) and administered 50 μl of 1 μg/μl ova-Alexa 647 (Invitrogen, Carlsbad, California) via oropharyngeal aspiration [42]. Lungs and MLNs were taken 2 and 48 hours post exposure and processed into single cell suspensions for flow analysis of CD11c<sup>+</sup> cells and uptake of ova-Alexa 647.

### Ex vivo co-culture of CD11c<sup>+</sup> pulmonary cells and CD4<sup>+</sup> T cells

CD11c<sup>+</sup> cells from the lung ( $2 \times 10^6$  cells/ml) were co-cultured with OTII CD4<sup>+</sup> T cells ( $1 \times 10^6$  cells/ml) and 100 μg/ml of ova<sub>323-339</sub> in a total volume of 200 μl/well in a flat-bottomed 96-well tissue culture plate. After 96 hours, cell-free supernatants were collected and frozen at -20°C. Cell-free supernatants were analyzed for the following mediators: IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, IL-12p40, IL-12p70, IL-13, IL-17, IFN $\gamma$ , IL-1 $\alpha$ , IL-1 $\beta$ , G-CSF, GM-CSF, KC, MCP-1, MIP-1 $\alpha$ , MIP-1 $\beta$ , TNF $\alpha$ , RANTES, and Eotaxin using Bio-Plex (Bio-Rad).


### Statistical analyses

Data were analyzed by two-tailed unpaired Student's *t* test or by two-way ANOVA with Bonferroni post test. Statistical calculations were performed using GraphPad Prism 5 for Windows. *p* values smaller than 0.05 were considered statistically significant.

### Results

#### Depletion of CD11c<sup>+</sup> cells prevents NO<sub>2</sub>-promoted allergic sensitization and manifestation of an allergic asthma phenotype

Optimal depletion of CD11c<sup>+</sup> cells in the spleen of CD11c-DTR mice has previously been reported to occur between 24 and 48 hours post i.p. administration of DT, with CD11c<sup>+</sup> cells beginning to re-populate by 72 hours [38]. To verify that depletion of CD11c<sup>+</sup> cells within the lung and MLN could be achieved in Tg<sup>+</sup> animals following intraperitoneal injection of DT, Tg<sup>-</sup> and Tg<sup>+</sup> mice were administered DT and then analyzed 24 hours later. Single-cell suspensions from the lung were stained for CD11c and analyzed by flow cytometry. CD11c<sup>+</sup> cells in the lung (Figure 1) were substantially depleted 24 hours



post administration of DT in CD11c-DTR Tg<sup>+</sup> mice. We next set out to test whether CD11c<sup>+</sup> cells were necessary during NO<sub>2</sub>-promoted allergic sensitization, and thus if depletion of these cells during sensitization would diminish features of allergic asthma in mice. We adjusted our previously-published model [6] to require only one sensitization, and thus only one administration of DT. Tg<sup>-</sup> and Tg<sup>+</sup> mice were administered DT via i.p. injection on day -1 and underwent inhalation of 15 ppm NO<sub>2</sub> for 1 hour followed by 30 minutes of aerosolized ova on day 0. Mice were then challenged with aerosolized ova on days 14, 15, and 16 and analyzed 48 hours later (day 18). Tg<sup>-</sup> mice exhibited features of allergic asthma, including elevated cell counts and eosinophils in BAL fluid, increased ova-specific IgE and IgG<sub>1</sub> in the serum, and secretion of Th2 and Th17 cytokines upon re-stimulation of CD4<sup>+</sup> T cells *in vitro*. Compared with Tg<sup>-</sup> mice, Tg<sup>+</sup> animals displayed significant decreases in the numbers of eosinophils and lymphocytes within the BAL (Figure 2A). In addition, when CD4<sup>+</sup> T cells were restimulated with APCs and ova, there were significant decreases in the levels of IL-4, IL-5, IL-13, and IL-17 (Figure 2B-E). Despite these decrease in cytokine production, there were no reductions in the serum levels of ova-specific IgE or IgG<sub>1</sub> (Figure 2F-G).

#### **NO<sub>2</sub> and ova exposure increases the number and maturation status of CD11c<sup>+</sup>CD11b<sup>-</sup> cells within the lung**


To understand immunological changes in the lung following NO<sub>2</sub> exposure that could allow for allergic sensitization in our mouse model [6], we investigated if the CD11c<sup>+</sup> cell populations within the lung were increased or activated following NO<sub>2</sub>/ova exposure. C57BL/6 mice were exposed to 1 hour of 15 ppm NO<sub>2</sub> followed by 30 minutes of aerosolized 3.4% ova and the lung and MLN were digested and analyzed by flow cytometry at 2 and 48 hours. Compared to air/ova exposed mice, bronchoalveolar lavage fluid recovered at 2 hours from the NO<sub>2</sub>/ova-exposed mice contained elevated levels of total protein (239.5 vs. 70.8 µg/ml; p = 0.040), IL-6 (315.8 vs. 157.5 pg/ml; p = 0.009), and MCP-1 (689.3 vs. 143.4 pg/ml; p = 0.026), indicative of the effects of 1 hour of 15 ppm NO<sub>2</sub> exposure, which manifest in a degree of lung damage and cytokine production similar to what we have previously reported [6]. Since pulmonary DCs are defined by a variety of cell surface markers, including primarily CD11c, CD11b, and MHCII [19,22,23], we assessed the number and maturation status within two separate cell populations that potentially contain DCs. These include CD11c<sup>+</sup>CD11b<sup>-</sup> cells and CD11c<sup>+</sup>CD11b<sup>+</sup> cells. While inflammatory CD11b<sup>+</sup>Gr-1<sup>lo</sup>F4/80<sup>lo</sup> DCs [22,27] were also studied, no significant alterations in this population were found and these data are therefore

not presented. 2 hours post NO<sub>2</sub>/ova exposure, the number of CD11c<sup>+</sup>CD11b<sup>-</sup> cells within the lung (Figure 3A) nearly doubled compared to the air/ova control (Figure 3B). Importantly, NO<sub>2</sub>-exposed CD11c<sup>+</sup>CD11b<sup>-</sup> cells expressed higher levels of MHCII, CD40, and OX40L (Figure 3C), indicating that resident CD11c<sup>+</sup> cells matured or that newly recruited CD11c<sup>+</sup> cells were of a more mature phenotype. These CD11c<sup>+</sup>CD11b<sup>-</sup> cells from air/ova- vs. NO<sub>2</sub>/ova-exposed lungs at 2 hours were on average 29.0 ± 1.5% vs. 42.4 ± 0.6% (p ≤ 0.01) MHCII<sup>+</sup>, 11.4 ± 0.09% vs. 16.3 ± 0.8% (p ≤ 0.05) CD40<sup>+</sup>, and 13.6 ± 1.2% vs. 16.3 ± 0.8% (p ≤ 0.05) OX40L<sup>+</sup>, differences that returned to baseline levels by 48 hours. CD11c<sup>+</sup> DCs expressing MHCII and CD40 have been shown to have a potent ability to cause T cell proliferation [16], while DCs expressing OX40L have been shown to induce naïve T cells to become Th2 polarized [43]. Thus, these changes in the CD11c<sup>+</sup>CD11b<sup>-</sup> cell population in the lung may have an important role in NO<sub>2</sub>-promoted allergic sensitization.

The number of lung cells within the CD11c<sup>+</sup>CD11b<sup>+</sup> population (Figure 4A) was not significantly altered following NO<sub>2</sub> exposure (Figure 4B). These cells were on average 81% ± 2.6% MHCII<sup>+</sup> regardless of air or NO<sub>2</sub> treatment or time point (data not shown). Despite already expressing high levels of MHCII, the expression level of the co-stimulatory molecules CD40 and CD86 did not change substantially following NO<sub>2</sub> exposure (Figure 4C). In fact, less than 10% of the CD11c<sup>+</sup>CD11b<sup>+</sup> population in the lung expressed CD40 and CD86 (data not shown). On average, 8.7% ± 2.1% of the cells were F4/80<sup>+</sup>, or potential macrophages [21]. CD11c and CD11b are typically expressed by myeloid DCs, the subset of DCs believed to be exceptionally well-suited for antigen presentation and activation of naïve CD4<sup>+</sup> T cells to become T effector cells [18]. Additionally, we measured an increase in CD11b expression within the CD11c<sup>+</sup>MHCII<sup>+</sup> population in the MLN following NO<sub>2</sub> exposure (Figure 5C), indicating that CD11c<sup>+</sup>CD11b<sup>+</sup> cells may be particularly important in trafficking to the lymph node and presenting antigen to T cells under pro-inflammatory conditions.


#### **Mature dendritic cells (CD11c<sup>+</sup>MHCII<sup>+</sup>) are present in the mediastinal lymph node (MLN) following NO<sub>2</sub> exposure**

The lung-draining mediastinal lymph nodes (MLNs) were pooled from 3 air/ova- or NO<sub>2</sub>/ova-exposed mice per group. Individual analysis of the BAL from each of these mice indicated they were all representative of their particular exposure group. In contrast to pulmonary macrophages, which are also CD11c<sup>+</sup>, pulmonary DCs migrate to draining lymph nodes upon maturation where they are able to stimulate naïve CD4<sup>+</sup> T cells



**Figure 2 Effects of depleting CD11c<sup>+</sup> cells during sensitization in an NO<sub>2</sub>-promoted allergic asthma model.** Eight- to fifteen-week-old female CD11c-DTR Tg<sup>+</sup> and Tg<sup>-</sup> mice were administered 4 ng DT/g of body weight via i.p. injection on day -1. All mice then underwent inhalation of 15 ppm NO<sub>2</sub> for 1 hour followed by 30 minutes of aerosolized ova on day 0. All mice were challenged with aerosolized ova on days 14, 15, and 16. Differential cell counts were measured from the BAL fluid 48 hours after the final ova challenge (A). Values shown are mean ± SEM with 8 Tg<sup>-</sup> and 5 Tg<sup>+</sup> mice per group. CD4<sup>+</sup> cells were isolated from spleens by positive selection on day 18 and co-cultured with antigen presenting cells (APCs) from naïve C57BL/6J mice and 100 µg/ml ova. Conditioned medium was collected at 96 hours and analyzed for the Th2 cytokines IL-4 (B), IL-5 (C), and IL-13 (D), and the Th17 cytokine IL-17 (E) by ELISA. No cytokines were detected in medium from APCs cultured alone or from CD4<sup>+</sup> T cells cultured with APCs in the absence of ova. Values shown are mean ± SEM with 4-5 mice per group. The ova-specific immunoglobulins IgE (F) and IgG<sub>1</sub> (G) were measured from serum collected 48 hours after the final ova challenge (day 18) by ELISA using serum from Alum/ova-immunized mice to generate standard curves. Values shown are mean ± SEM with 7-10 mice per group. Statistics were computed by unpaired Student's t test. \* denotes *p* < 0.05 and \*\* denotes *p* < 0.01 compared with respective Tg<sup>-</sup> samples.





**Figure 3 Number and maturation status of lung CD11c<sup>+</sup>CD11b<sup>-</sup> cells following NO<sub>2</sub> and ova exposure.** Mice were exposed to 1 hour of 15ppm NO<sub>2</sub> or air followed by 30 minutes of aerosolized 3.4% ova. Lungs were harvested 2 and 48 hours later, single cell suspensions were generated and stained with antibodies, and cells were analyzed by flow cytometry. Dead cells were excluded by FSC and SSC gating and then total lung cells were gated comparing CD11c and CD11b expression, with future analyses focusing on the CD11c<sup>+</sup>CD11b<sup>-</sup> population (bold gate, A). The total number of CD11c<sup>+</sup>CD11b<sup>-</sup> cells within the lung was enumerated (B) and the maturation status of these cells was assessed by median fluorescence intensity (MFI) of MHCII and the co-stimulatory molecules CD40 and OX40L 2 hours post exposure (C). NO<sub>2</sub>-exposed animals are represented by a thick line or black bar while air-exposed animals are represented by a thin line or white bar. Grey, filled histograms are isotype controls. Data shown are mean ± SEM with 3 animals per group and are representative of experiments performed twice. \* denotes  $p < 0.05$  and \*\* denotes  $p < 0.01$  versus air control at the same time point by unpaired Student's t test. ^ denotes  $p < 0.05$  and ^^ denotes  $p < 0.01$  versus 2 hours post NO<sub>2</sub> exposure by unpaired Student's t test.


[13]. 2 hours following exposure to 15 ppm NO<sub>2</sub> and 3.4% ova, there was a slight increase in CD11c<sup>+</sup>MHCII<sup>+</sup> DCs within the MLN (3.71% to 4.17% of CD45<sup>+</sup> cells) (Figure 5A). At 2 hours, these CD11c<sup>+</sup>MHCII<sup>+</sup> cells displayed markers indicative of maturation (Figure 5B). However, by 48 hours post exposure to NO<sub>2</sub> and ova, MLN DCs displayed dramatic upregulation of the co-stimulatory molecules CD80, CD86, and OX40L, consistent with maturation and a phenotype capable of inducing a T cell-mediated inflammatory response

[16,29] (Figure 5C). Furthermore, infiltrating MLN DCs were increasingly CD11b<sup>+</sup> at 48 hours, consistent with the phenotype of a myeloid or inflammatory DC.

#### NO<sub>2</sub>-exposed pulmonary CD11c<sup>+</sup> cells produce pro-inflammatory cytokines and induce T cell cytokine production ex vivo

Mature DCs have been documented to possess increased antigen presentation capabilities, strongly inducing T cell proliferation and cytokine production [16].






**Figure 4** Number and maturation status of lung CD11c<sup>+</sup>CD11b<sup>+</sup> cells following NO<sub>2</sub> and ova exposure. Mice were exposed to 1 hour of 15 ppm NO<sub>2</sub> or Air followed by 30 minutes of aerosolized 3.4% ova. Lungs were harvested 2 and 48 hours later, single cell suspensions were generated and stained with antibodies, and cells were analyzed by flow cytometry. Dead cells were excluded by FSC and SSC gating and then total lung cells were gated comparing CD11c and CD11b expression, with future analyses focusing on the CD11c<sup>+</sup>CD11b<sup>+</sup> population (bold gate, A). The total number of CD11c<sup>+</sup>CD11b<sup>+</sup> cells within the lung was enumerated (B) and the maturation status of these cells was assessed by expression of MHCII and the co-stimulatory molecules CD40 and CD86 48 hours post exposure (C). Maturation markers are graphed as median fluorescence intensity (MFI). NO<sub>2</sub>-exposed animals are represented by a thick line or black bar while air-exposed animals are represented by a thin line or white bar. Grey, filled histograms are isotype controls. Data shown are mean ± SEM with 3 animals per group and are representative of experiments performed twice. Differences did not reach statistical significance by unpaired Student's t test.

Therefore, we investigated whether CD11c<sup>+</sup> pulmonary cells from NO<sub>2</sub>-exposed animals have increased pro-inflammatory and T cell activating properties compared to those from air-exposed mice. Mice were exposed to either air or 15 ppm NO<sub>2</sub> for one hour followed by 30 minutes of 3.4% ova. We then waited 48 hours as we have previously reported that separating the time

between NO<sub>2</sub> and the initial ova exposure increases sensitization to ova [6]. Thus, 48 hours following NO<sub>2</sub> and ova exposure, CD11c<sup>+</sup> cells were removed from total lung digestions by positive selection and co-cultured with naïve splenic CD4<sup>+</sup> transgenic OTII T cells, which possess T cell receptors specific only for the antigenic peptide of ovalbumin, ova<sub>323-339</sub> [39]. Since the majority





**Figure 5** CD11c<sup>+</sup> dendritic cell maturation status in mediastinal lymph nodes (MLN) following NO<sub>2</sub> and ova exposure. Mice were exposed to 1 hour of 15 ppm NO<sub>2</sub> or air followed by 30 minutes of aerosolized 3.4% ova. Lungs were harvested 2 and 48 hours later, single cell suspensions were generated and stained with antibodies, and cells were analyzed by flow cytometry. Only CD45<sup>+</sup> MLN cells were included in analyses. NO<sub>2</sub>-exposed animals are represented by a thick line while air-exposed animals are represented by a thin line. Grey, filled histograms are isotype controls. CD11c<sup>+</sup>MHCII<sup>+</sup> cells were measured as a percent of all CD45<sup>+</sup> MLN cells (A) and were analyzed for expression of the markers CD80, CD86, CD40, OX40L, and CD11b at 2 hours (B) and 48 hours (C) post exposure. Marker expression is shown as median fluorescence intensity (MFI). Data shown represent MLN cells pooled from 3 animals per group and are representative of experiments performed twice.

of the immunostimulatory CD11c<sup>+</sup> cells that had engulfed the ovalbumin encountered during the inhaled exposure were likely no longer in the lung (see figures 6C and 7C), the cells were then either cultured *in vitro* with 100 µg of ova<sub>323-339</sub> (stimulated) or in media alone (unstimulated). The cells were co-cultured for 96 hours and the supernatants were then analyzed for an array of cytokines and chemokines associated with DC activation, T cell activation, and Th2 polarization. When comparing stimulated cells, the CD11c<sup>+</sup> cells from NO<sub>2</sub>-exposed animals produced significant amounts of IL-1α and IL-1β, a Th17-polarizing cytokine [34], versus air-exposed controls (Figure 8A). NO<sub>2</sub>-exposed CD11c<sup>+</sup> cells also secreted increased quantities of IL-12, a Th1 polarizing cytokine, and IL-6, a Th2 polarizing cytokine [30,31](Figure 8B). Interestingly, air-exposed and NO<sub>2</sub>-exposed CD11c<sup>+</sup> cells were equally capable of promoting T cell proliferation, as measured by IL-2 production (Figure 8C). However, the T cells co-cultured with NO<sub>2</sub>-exposed CD11c<sup>+</sup> cells produced significantly larger amounts of IL-5 (Figure 8C), with no difference in IL-13 or IL-17 (Figure 8D).

#### NO<sub>2</sub> induces pulmonary CD11c<sup>+</sup> cells to capture antigen and travel to the mediastinal lymph node

DCs are particularly well-equipped for activating T cells as they are capable of capturing and processing antigens, travelling to the draining lymph node, and presenting those antigens to T cells [16,18]. Under inflammatory conditions, DCs increase in rate of turnover and total numbers within the lung and in transit to the draining lymph node [44,45]. To further study DC trafficking and maturation following NO<sub>2</sub> exposure, Alexa 647-labeled ova was employed [21]. To test if conventional CD11c<sup>+</sup> DCs increase antigen uptake following NO<sub>2</sub> exposure, C57BL/6 mice were exposed to 1 hour of 15 ppm NO<sub>2</sub> or air followed by oropharyngeal aspiration of ova-Alexa 647. We focused on cells expressing CD11c as these populations displayed maturation following NO<sub>2</sub> exposure in previous studies (Figure 3). We separately analyzed CD11c<sup>+</sup>CD11b<sup>-</sup> and CD11c<sup>+</sup>CD11b<sup>+</sup> cells, as before, to identify differences between these two populations that contain DCs (Figure 6A). Two hours post-exposure, pulmonary CD11c<sup>+</sup>CD11b<sup>-</sup> cells contained increased amounts of ova-Alexa 647 in the NO<sub>2</sub>- versus air-exposed mice (Figure 6C). At 48 hours, ova-Alexa 647<sup>+</sup> CD11c<sup>+</sup>CD11b<sup>-</sup> cells were still present in both the air- and NO<sub>2</sub>-treated groups with equivalent retention of labeled ova per cell (Figure 6C) despite the increased percentage of CD11c<sup>+</sup>CD11b<sup>-</sup> cells in the lungs of NO<sub>2</sub>-exposed animals (Figure 6B).






**Figure 7 Pro-inflammatory cytokine production and induction of CD4<sup>+</sup> T cell cytokine production ex vivo by NO<sub>2</sub>-exposed pulmonary CD11c<sup>+</sup> cells.** Mice were exposed to air or 15 ppm NO<sub>2</sub> for 1 hour followed by 30 minutes of aerosolized 3.4% ova. Forty-eight hours later, CD11c<sup>+</sup> cells were purified from lungs via positive selection and used as antigen presenting cells co-cultured with CD4<sup>+</sup> T cells purified from the spleen of OTII transgenic mice via positive selection. CD11c<sup>+</sup> cells (2 × 10<sup>6</sup> cells/ml) co-cultured with T cells (1 × 10<sup>6</sup> cells/ml) were stimulated with 100 μg/ml ova<sub>323-339</sub> and cultured for 4 days or left unstimulated in media alone. Conditioned medium were analyzed via BioPlex for pro-inflammatory mediators associated with CD11c<sup>+</sup> cell activation (IL-1α and IL-1β) (A) and T cell polarization (IL-12p70 and IL-6) (B). Cytokines associated with T cell proliferation (IL-2) (C) as well as cytokines produced by activated Th2 cells (IL-5 and IL-13) (C, D) and Th17 cells (IL-17) (D) were measured. Data shown are mean ± SEM with 6 mice per group. \* denotes p < 0.05, \*\* denotes p < 0.01, \*\*\* denotes p < 0.001 compared with Air Stim as determined by one-way ANOVA with Bonferroni post test.

Furthermore, while the total cell numbers from the lung were unchanged by NO<sub>2</sub>/ova, CD11c<sup>+</sup>CD11b<sup>+</sup> cells increased as a percentage of total lung cells 2 hours post NO<sub>2</sub>/ova-Alexa 647 exposure versus air control and elevated further by 48 hours (Figure 7B). NO<sub>2</sub> exposure induced CD11c<sup>+</sup>CD11b<sup>+</sup> cells to increase antigen capture, as shown by a shift in the median fluorescence intensity of ova-Alexa 647 at 2 hours compared with the air control (Figure 7C). These CD11c<sup>+</sup>CD11b<sup>+</sup>ova-Alexa 647<sup>+</sup> cells were diminished at 48 hours versus the air control, potentially due to migration of this population to the MLN (Figure 7C, 9B, C). As shown previously, CD11c<sup>+</sup>CD11b<sup>+</sup> cells from both air- and NO<sub>2</sub>-exposed animals expressed high levels of MHCII (Figure 4C), but the levels of MHCII were significantly higher on CD11c<sup>+</sup>MHCII<sup>+</sup> cells from NO<sub>2</sub>-exposed MLNs at 2 hours (Figure 9D). By 48 hours, CD11c<sup>+</sup>MHCII<sup>+</sup> cells (DCs) in the MLN contained increased amounts of ova-Alexa 647 in NO<sub>2</sub>- versus air-exposed animals (Figure 9B, C), indicating that NO<sub>2</sub> exposure promoted antigen capture and DC transit to the draining lymph node.

## Discussion

The incidence of allergic asthma continues to rise throughout industrialized countries and nations in transition [1], where the concentrations of air pollutants, such as NO<sub>2</sub>, are also increasing [46,47]. Studies have correlated an increased likelihood of developing asthma [12] and worsening asthma symptoms [10] with high levels of ambient NO<sub>2</sub>. Respiratory viral infections also result in NO<sub>2</sub> production within the lung [48,49] and data suggest that certain severe viral infections early in life increase the risk of developing asthma [48]. In our previous studies, we have demonstrated that in mice, NO<sub>2</sub> can act as an adjuvant, promoting allergic sensitization to an innocuous inhaled antigen [6]. Therefore, understanding the mechanisms by which NO<sub>2</sub> promotes inappropriate adaptive immune responses will provide powerful insight concerning the effects of pollution and viral infection on respiratory health, the development of asthma, the balance between inhalational tolerance and allergy, and potentially provide mechanistic targets for prevention or treatment.




**Figure 8** Effects of NO<sub>2</sub> inhalation on antigen capture in CD11c<sup>+</sup>CD11b<sup>-</sup> lung cells *in vivo*. Mice were exposed to air or 15 ppm NO<sub>2</sub> for one hour followed by oropharyngeal aspiration of 50 μg of ova-Alexa 647. Lungs were harvested 2 and 48 hours later, single cell suspensions were generated and stained with antibodies, and cells were analyzed by flow cytometry. Dead cells were excluded by FSC and SSC gating and then total lung cells were gated comparing CD11c and CD11b expression, with future analyses focusing on the CD11c<sup>+</sup>CD11b<sup>-</sup> population (bold gate, A). The percent of CD11c<sup>+</sup>CD11b<sup>-</sup> cells within the lung was determined (B) and uptake of the antigen ova-Alexa 647 was quantified by median fluorescence intensity (MFI) (C). NO<sub>2</sub>-exposed animals are represented by a thick line while air-exposed animals are represented by a thin line. Grey, filled histograms are naive controls (did not receive ova-Alexa 647) (C). Data shown are mean ± SEM with 4 animals per group and are representative of experiments performed twice. \* denotes  $p < 0.05$  and \*\* denotes  $p < 0.01$  versus air control at the same time point by unpaired Student's t test. ^^ denotes  $p < 0.01$  and ^^ denotes  $p < 0.001$  versus 2 hours post NO<sub>2</sub> exposure by unpaired Student's t test.

CD11c<sup>+</sup> DCs are critical in activating naïve CD4<sup>+</sup> T cells [16], the products of which are primarily responsible for the symptoms of allergic asthma [36,50]. Multiple inhaled environmental agents have adjuvant activities and lead to allergic sensitization, including endotoxin [51,52], triacylated lipopeptides [53], residual oil fly ash [54,55], diesel exhaust particles [56-58], cigarette smoke [59], ultrafine particles [60], and house dust mite [61,62]. Proposed mechanisms for this inappropriate immune response promoted by environmental agents include disruption of the epithelial barrier, leading to cytokine and chemokine production by epithelial cells, as well as antigen access to underlying DCs [25,63] and altered DC function [60,64-69]. Because CD11c<sup>+</sup> DCs are crucial for mounting naïve T cell responses, we hypothesized that eliminating CD11c<sup>+</sup> cells during NO<sub>2</sub>-promoted allergic sensitization would minimize features of allergic asthma in mice. To test this hypothesis, we utilized transgenic CD11c-DTR mice, in which temporary depletion of CD11c<sup>+</sup> DCs is possible.

To use the CD11c-DTR mice, we adjusted our model of NO<sub>2</sub>-promoted allergic sensitization to include only

one sensitization. Our data demonstrate that even a single exposure to 15 ppm NO<sub>2</sub> for 1 hour followed by 30 minutes of aerosolized ova immediately after is sufficient to induce allergic sensitization to ova. Interestingly, we measured substantial increases in IL-4, IL-5, IL-13, and IL-17 from restimulated CD4<sup>+</sup> T cells as well as substantial increases in eosinophilia and neutrophilia in the BAL, indicating that our one sensitization model polarizes CD4<sup>+</sup> T cells down both Th2 and Th17 pathways. These pathways overlap as the mediator IL-6 is important in promoting both pathways [30-32] and was secreted at elevated levels by DCs from NO<sub>2</sub>-exposed mice. Systemic administration of DT to CD11c-DTR transgenic mice prior to sensitization resulted in a decrease in the total white blood cell count and eosinophilia in the BAL of Tg<sup>+</sup> animals versus Tg<sup>-</sup> animals. Similar differences were measured for cytokine production from *in vitro* restimulated CD4<sup>+</sup> T cells. Thus, systemic depletion of CD11c<sup>+</sup> cells provides mechanistic insight into the important role of CD11c<sup>+</sup> cells in NO<sub>2</sub>-promoted allergic sensitization. However, our findings suggest that while CD11c<sup>+</sup> cells



**Figure 9 Effects of NO<sub>2</sub> inhalation on antigen-containing dendritic cells in the mediastinal lymph node.** Mice were exposed to air or 15 ppm NO<sub>2</sub> for 1 hour followed by oropharyngeal aspiration of 50 µg of ova-Alexa 647. MLN were harvested 2 and 48 hours later, single cell suspensions were generated and stained with antibodies, and cells were analyzed by flow cytometry. CD11c<sup>+</sup>MHCII<sup>+</sup> cells (DCs) were gated as a percentage of CD45<sup>+</sup> cells in the MLN (A). DC retention of the antigen ova-Alexa 647 was quantified by the median fluorescence intensity (MFI) of Alexa 647 (B). The histogram represents ova-Alexa 647 retention at 48 hours post exposure (C). NO<sub>2</sub>-exposed animals are represented by a thick line while air-exposed animals are represented by a thin line. Grey, filled histograms are naive controls (did not receive ova-Alexa 647) (C). The maturation status of the DC population was assessed by expression of MHCII (D). Data shown are mean ± SEM with 4 animals per group and are representative of experiments performed twice. \* denotes *p* < 0.05 versus air control at the same time point by unpaired Student's *t* test. ^ denotes *p* < 0.05 versus 2 hours post NO<sub>2</sub> exposure by unpaired Student's *t* test.

are important in NO<sub>2</sub>-promoted allergic sensitization, depleting these cells during sensitization does not abolish all parameters of the allergic asthma phenotype, as we were unable to measure a difference in the levels of Tg<sup>+</sup> and Tg<sup>-</sup> animals. This may be due, in part, to the low levels of antigen-specific immunoglobulins generated as a consequence of our modified sensitization regimen in which mice were only exposed to inhaled NO<sub>2</sub> and ova on a single occasion compared to our previously-published model that involved two sensitizations [6]. Importantly, when DT was administered via i.p. injection, depletion of CD11c<sup>+</sup> cells within the lung occurred as expected in Tg<sup>+</sup> animals [17]. However, this depletion does leave some CD11c<sup>+</sup> cells present in the lung, inducing approximately an 80% decrease from 5.8% to 1% of total lung cells.

Thus, it is possible that this depletion minimizes the initial number of naïve T cells activated, but that after repeated exposure to antigen, clonal expansion allows for the development of some of the features of the Th2 response, including the expansion of the eosinophil population, their recruitment to the airway, and the generation of antigen-specific immunoglobulins. Additionally, depleting CD11c<sup>+</sup> cells during sensitization may alter rather than reduce the immune response, changing the outcomes associated with sensitization. However, we did not measure any qualitative alterations in the type of immune response provoked in the CD11c-DTR Tg<sup>+</sup> mice.

It is important to consider other CD11c<sup>+</sup> cell populations in addition to mDCs that are also affected by DT administration to Tg<sup>+</sup> animals. Plasmacytoid DCs (pDCs) and alveolar macrophages also express CD11c,



albeit to a lesser extent than mDCs [21]. Plasmacytoid DCs promote tolerance by decreasing the capability of myeloid DCs to induce T cell proliferation as well as promoting CD4<sup>+</sup> T regulatory cell proliferation [26]. Thus, by depleting CD11c<sup>+</sup> cells within the lung, depletion of both pro-inflammatory mDCs as well as anti-inflammatory pDCs occurs. Despite this caveat, CD11c-DTR mice have revealed the importance of CD11c<sup>+</sup> cells in promoting allergic sensitization in other models.

Our current studies demonstrate that one mechanism by which NO<sub>2</sub> may act as an adjuvant is through the activation and recruitment of pulmonary CD11c<sup>+</sup> DCs, a cell type believed to act as the bridge between innate and adaptive immunity [25]. We measured increased numbers of CD11c<sup>+</sup>CD11b<sup>-</sup> cells and minimal changes in CD11c<sup>+</sup>CD11b<sup>+</sup> cells in the lungs 2 hours following NO<sub>2</sub> exposure, all of which have the potential to be DCs. Other adjuvants have been shown to work similarly, including the well known adjuvant alum, which causes the recruitment of inflammatory DCs to the site of injection (peritoneum) and subsequently to the draining lymph nodes [22]. Additionally, the CD11c<sup>+</sup>CD11b<sup>-</sup> population in the lung significantly upregulated expression of MHCII, CD40, and OX40L 2 hours post-NO<sub>2</sub>, thereby enhancing its capability for antigen presentation and providing co-stimulation to promote Th2 responses. Myeloid DCs pulsed with ova and administered directly into mouse lungs also result in allergic sensitization to ova, indicating that these cells have the capability to induce T cell responses [24]. Furthermore, our data demonstrate that inhalation of 15 ppm NO<sub>2</sub> for just 1 hour is sufficient to induce the production of MCP-1 within the lung, a chemokine associated with DC recruitment [70]. IL-6 is also increased in the BAL fluid 2 hours following NO<sub>2</sub> exposure, which can induce a Th2 polarized environment [31].

From NO<sub>2</sub>-exposed mice, we measured increased protein in the BAL, indicative of cellular damage that may promote the activation of epithelial cells capable of secreting factors such as IL-6 [71] and other pro-inflammatory mediators [25]. Agents that cause damage to the lung epithelium, such as the proteases produced by house dust mites, have been shown to promote allergic sensitization through allowing access of the antigens to resident antigen presenting cells and inducing cytokine and chemokine production from epithelial cells [63]. Previous findings have demonstrated that 10 ppm NO<sub>2</sub> induces the release of intracellular molecules such as HSP70 [6], which can act as an endogenous danger signal [72] and activate the transcription factor NF- $\kappa$ B, a key modulator in both innate and adaptive immune responses. Interestingly, NF- $\kappa$ B is capable of inducing IL-6 and MCP-1 gene expression [73] and is activated following NO<sub>2</sub> exposure [6], providing a potential mechanism by which NO<sub>2</sub> promotes CD11c<sup>+</sup> pulmonary

cell activation and Th2 polarization. Our *in vitro* co-culture data indicate that an effect of NO<sub>2</sub> solely on lung DCs is insufficient to explain the creation of a Th2/Th17-polarizing environment in the lung. *In vivo*, additional cell types, including epithelium, contribute to the effects of NO<sub>2</sub> inhalation on CD4<sup>+</sup> T cell polarization. We have recently reported that NF- $\kappa$ B activation in the airway epithelium, as is also induced by NO<sub>2</sub> exposure [6,74], is sufficient to activate mediastinal lymph node DCs and to promote a mixed Th2/Th17 response to an innocuous inhaled antigen [75].

Allergic asthma is primarily dependent upon the activities of CD4<sup>+</sup> T cells [2]. CD11c<sup>+</sup> DCs regulate the initiation of naive CD4<sup>+</sup> T cell responses in multiple ways, including carrying antigens to the draining lymph node, processing and presenting these antigens, providing co-stimulation, and secreting cytokines [16]. In our studies, CD11c<sup>+</sup> pulmonary cells secreted increased amounts of IL-1 $\alpha$ , IL-1 $\beta$ , IL-12p70 and IL-6 *ex vivo* following NO<sub>2</sub> exposure, demonstrating that these cells alone produce pro-inflammatory mediators. When co-cultured with CD4<sup>+</sup> OTII cells, NO<sub>2</sub>-exposed pulmonary CD11c<sup>+</sup> cells promoted the production of IL-5 from OTII cells more so than air-exposed controls, implicating that NO<sub>2</sub>-exposed CD11c<sup>+</sup> cells induced increased Th2 cell activity. Although we measured increases in Th1- (IL-12p70), Th2- (IL-6), and Th17-(IL-1) polarizing cytokines, it is important to remember that these co-cultures did not include epithelial cells. It is likely that epithelial cells also influence the outcome of DC and CD4<sup>+</sup> T cell interactions, as epithelial cells are capable of secreting multiple mediators that affect T cell polarization, promoting the development of a Th2 immune response [25]. This may also be one of the reasons why we observed no differences in the secretion of IL-17 by the CD4<sup>+</sup> T cells co-cultured with DCs from NO<sub>2</sub>-exposed mice compared to those from the air-exposed mice.

NO<sub>2</sub> exposure also caused pulmonary CD11c<sup>+</sup>CD11b<sup>-</sup> and CD11c<sup>+</sup>CD11b<sup>+</sup> cells to increase antigen capture and trafficking to the draining lymph node, increasing the opportunity for a T cell response. Interestingly, 2 hours following NO<sub>2</sub>, both CD11c<sup>+</sup>CD11b<sup>-</sup> and CD11c<sup>+</sup>CD11b<sup>+</sup> cells within the lung contained increased amounts of ova-Alexa 647, but by 48 hours contained equal (CD11c<sup>+</sup>CD11b<sup>-</sup>) or lower (CD11c<sup>+</sup>CD11b<sup>+</sup>) amounts of ova-Alexa 647 than air controls. Also at 48 hours, CD11c<sup>+</sup>MHCII<sup>+</sup> cells containing ova-Alexa 647 increased in the MLN, indicating that NO<sub>2</sub> inhalation induced trafficking of these cells to the MLN.

## Conclusions

Collectively, the results of the studies presented herein demonstrate that CD11c<sup>+</sup> cells, which include conventional DCs, are critical in NO<sub>2</sub>-promoted allergic



sensitization, as depletion of these cells only during sensitization diminished multiple features of allergic asthma in mice. NO<sub>2</sub> inhalation significantly impacts pulmonary CD11c<sup>+</sup> cells, as shown by increased cytokine production, upregulation of maturation markers, increased antigen uptake, and improved ability to stimulate naïve T cells, all of which occur in a temporally-coordinated manner. Future work will be required to identify critical mediators of these effects through the use of neutralizing antibodies and knockout mice, as well as by further refining the direct or indirect effects of NO<sub>2</sub> on specific subsets of pulmonary DCs, including the CD103<sup>+</sup> population that expresses adherens junction proteins and interacts closely with the epithelial cells at the interface of the airway with the external environment [76]. Additionally, we currently cannot exclude the possibility that other factors affected by NO<sub>2</sub> exposure, such as the pH of the airway surface lining fluid and oxidation/nitration events within proteins [77-79], may play a role in the adjuvant effects of NO<sub>2</sub>. Understanding the mechanisms underlying the inappropriate T cell activation induced following exposure to environmental pollutants or endogenously-generated oxidant gasses, such as NO<sub>2</sub>, will provide insight to allergic sensitization as it occurs in humans, providing further understanding of the causes of allergic asthma.

#### List of Abbreviations

APC: antigen-presenting cell; BAL: bronchoalveolar lavage; CD: cluster of differentiation; DC: dendritic cell; DT: diphtheria toxin; DTR: diphtheria toxin receptor; IL: interleukin; MHCII: major histocompatibility complex class 2; MLN: mediastinal lymph node; NO<sub>2</sub>: nitrogen dioxide; ova: ovalbumin; ppm: parts per million; Tg+: transgenic; Tg-: transgene negative; Th: T helper.

#### Acknowledgements

The authors thank Dr. Karen Fortner (UVM College of Medicine, Immunobiology) and Colette Charland, Director of the University of Vermont's Immunobiology Flow Cytometry Core, for helpful discussions and reagent recommendations. This work was supported by R01 HL089177 (MEP), NCRN COBRE P20RR15557 (LAWL), R01 HL084200 (BTS), and R01 AI067897 (JEB).

#### Author details

<sup>1</sup>The Vermont Lung Center and Department of Medicine, University of Vermont, Burlington, VT, 05405, USA. <sup>2</sup>Department of Surgery, University of Vermont, Burlington, VT, 05405, USA.

#### Authors' contributions

SRH designed the experiments, executed the studies, and drafted the manuscript. JLA bred, genotyped, and maintained mice for the studies. SAP and JLA assisted with cell staining for flow cytometry. LAWL assisted with study design and cell staining. BTS and JEB assisted with analysis of flow cytometry data. MEP conceived of the studies, coordinated their execution, and revised the manuscript. All authors read and approved the final manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 15 March 2010 Accepted: 26 July 2010  
Published: 26 July 2010

#### References

1. Mannino DM, Homa DM, Pertowski CA, Ashizawa A, Nixon LL, Johnson CA, Ball LB, Jack E, Kang DS: **Surveillance for asthma—United States, 1960-1995.** *MMWR CDC Surveill Summ* 1998, **47**(1):1-27.
2. Herrick CA, Bottomly K: **To respond or not to respond: T cells in allergic asthma.** *Nat Rev Immunol* 2003, **3**(5):405-412.
3. Al-Ramli W, Prefontaine D, Chouiali F, Martin JG, Olivenstein R, Lemiere C, Hamid Q: **T(H)17-associated cytokines (IL-17A and IL-17F) in severe asthma.** *J Allergy Clin Immunol* 2009, **123**(5):1185-1187.
4. McKinley L, Alcorn JF, Peterson A, Dupont RB, Kapadia S, Logar A, Henry A, Irvin CG, Piganelli JD, Ray A, et al: **TH17 cells mediate steroid-resistant airway inflammation and airway hyperresponsiveness in mice.** *J Immunol* 2008, **181**(6):4089-4097.
5. Alcorn JF, Crowe CR, Kolls JK: **TH17 cells in asthma and COPD.** *Annu Rev Physiol* 2010, **72**:495-516.
6. Bevelander M, Mayette J, Whittaker LA, Paveglia SA, Jones CC, Robbins J, Hemenway D, Akira S, Uematsu S, Poynter ME: **Nitrogen dioxide promotes allergic sensitization to inhaled antigen.** *J Immunol* 2007, **179**(6):3680-3688.
7. Dykewicz MS: **Occupational asthma: current concepts in pathogenesis, diagnosis, and management.** *J Allergy Clin Immunol* 2009, **123**(3):519-528.
8. Beelen R, Hoek G, van den Brandt PA, Goldbohm RA, Fischer P, Schouten LJ, Jerrett M, Hughes E, Armstrong B, Brunekreef B: **Long-term effects of traffic-related air pollution on mortality in a Dutch cohort (NLCS-AIR study).** *Environ Health Perspect* 2008, **116**(2):196-202.
9. Poynter ME, Persinger RL, Irvin CG, Butnor KJ, van Hirtum H, Blay W, Heintz NH, Robbins J, Hemenway D, Taatjes DJ, et al: **Nitrogen dioxide enhances allergic airway inflammation and hyperresponsiveness in the mouse.** *Am J Physiol Lung Cell Mol Physiol* 2006, **290**(1):L144-152.
10. Kattan M, Gergen PJ, Eggleston P, Visness CM, Mitchell HE: **Health effects of indoor nitrogen dioxide and passive smoking on urban asthmatic children.** *J Allergy Clin Immunol* 2007, **120**(3):618-624.
11. Pattenden S, Hoek G, Braun-Fahrländer C, Forastiere F, Kosheleva A, Neuberger M, Fletcher T: **NO<sub>2</sub> and children's respiratory symptoms in the PATY study.** *Occup Environ Med* 2006, **63**(12):828-835.
12. Gauderman WJ, Avol E, Lurmann F, Kuenzli N, Gilliland F, Peters J, McConnell R: **Childhood asthma and exposure to traffic and nitrogen dioxide.** *Epidemiology* 2005, **16**(6):737-743.
13. Vermaelen KY, Carro-Muino I, Lambrecht BN, Pauwels RA: **Specific migratory dendritic cells rapidly transport antigen from the airways to the thoracic lymph nodes.** *J Exp Med* 2001, **193**(1):51-60.
14. Vermaelen K, Pauwels R: **Pulmonary dendritic cells.** *Am J Respir Crit Care Med* 2005, **172**(5):530-551.
15. Holt PG, Strickland DH, Wikstrom ME, Jahnsen FL: **Regulation of immunological homeostasis in the respiratory tract.** *Nat Rev Immunol* 2008, **8**(2):142-152.
16. Bancheau J, Steinman RM: **Dendritic cells and the control of immunity.** *Nature* 1998, **392**(6673):245-252.
17. van Rijst LS, Jung S, Kleinjan A, Vos N, Willart M, Duez C, Hoogsteden HC, Lambrecht BN: **In vivo depletion of lung CD11c<sup>+</sup> dendritic cells during allergen challenge abrogates the characteristic features of asthma.** *J Exp Med* 2005, **201**(6):981-991.
18. Shortman K, Liu YJ: **Mouse and human dendritic cell subtypes.** *Nat Rev Immunol* 2002, **2**(3):151-161.
19. Lewkowich IP, Lajoie S, Clark JR, Herman NS, Sproles AA, Wills-Karp M: **Allergen uptake, activation, and IL-23 production by pulmonary myeloid DCs drives airway hyperresponsiveness in asthma-susceptible mice.** *PLoS ONE* 2008, **3**(12):e3879.
20. Akdis M, Verhagen J, Taylor A, Karamloo F, Karagiannidis C, Cramer R, Thunberg S, Deniz G, Valenta R, Fiebigh H, et al: **Immune responses in healthy and allergic individuals are characterized by a fine balance between allergen-specific T regulatory 1 and T helper 2 cells.** *J Exp Med* 2004, **199**(11):1567-1575.
21. Wikstrom ME, Stumbles PA: **Mouse respiratory tract dendritic cell subsets and the immunological fate of inhaled antigens.** *Immunol Cell Biol* 2007, **85**(3):182-188.
22. Kool M, Soullie T, van Nimwegen M, Willart MA, Muskens F, Jung S, Hoogsteden HC, Hammad H, Lambrecht BN: **Alum adjuvant boosts adaptive immunity by inducing uric acid and activating inflammatory dendritic cells.** *J Exp Med* 2008, **205**(4):869-882.

23. Vermaelen K, Pauwels R: **Accurate and simple discrimination of mouse pulmonary dendritic cell and macrophage populations by flow cytometry: methodology and new insights.** *Cytometry A* 2004, **61**(2):170-177.
24. Lambrecht BN, De Veerman M, Coyle AJ, Gutierrez-Ramos JC, Thielemans K, Pauwels RA: **Myeloid dendritic cells induce Th2 responses to inhaled antigen, leading to eosinophilic airway inflammation.** *J Clin Invest* 2000, **106**(4):551-559.
25. Hammad H, Lambrecht BN: **Dendritic cells and epithelial cells: linking innate and adaptive immunity in asthma.** *Nat Rev Immunol* 2008, **8**(3):193-204.
26. de Heer HJ, Hammad H, Soullie T, Hijdra D, Vos N, Willart MA, Hoogsteden HC, Lambrecht BN: **Essential role of lung plasmacytoid dendritic cells in preventing asthmatic reactions to harmless inhaled antigen.** *J Exp Med* 2004, **200**(1):89-98.
27. Randolph GJ, Inaba K, Robbiani DF, Steinman RM, Muller WA: **Differentiation of phagocytic monocytes into lymph node dendritic cells in vivo.** *Immunity* 1999, **11**(6):753-761.
28. Deurloo DT, van Berkel MA, van Esch BC, Hofhuis F, Nijkamp FP, Oosterwegel MA, van Oosterhout AJ: **CD28/CTLA4 double deficient mice demonstrate crucial role for B7 co-stimulation in the induction of allergic lower airways disease.** *Clin Exp Allergy* 2003, **33**(9):1297-1304.
29. Hoshino A, Tanaka Y, Akiba H, Asakura Y, Mita Y, Sakurai T, Takaoka A, Nakaïke S, Ishii N, Sugamura K, et al: **Critical role for OX40 ligand in the development of pathogenic Th2 cells in a murine model of asthma.** *Eur J Immunol* 2003, **33**(4):861-869.
30. Krishnamoorthy N, Oriss TB, Paglia M, Fei M, Yarlagadda M, Vanhaesebroeck B, Ray A, Ray P: **Activation of c-Kit in dendritic cells regulates T helper cell differentiation and allergic asthma.** *Nat Med* 2008, **14**(5):565-573.
31. Rincon M, Anguita J, Nakamura T, Fikrig E, Flavell RA: **Interleukin (IL)-6 directs the differentiation of IL-4-producing CD4+ T cells.** *J Exp Med* 1997, **185**(3):461-469.
32. Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, Weiner HL, Kuchroo VK: **Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells.** *Nature* 2006, **441**(7090):235-238.
33. Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B: **TGFbeta in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells.** *Immunity* 2006, **24**(2):179-189.
34. Sutton C, Brereton C, Keogh B, Mills KH, Lavelle EC: **A crucial role for interleukin (IL)-1 in the induction of IL-17-producing T cells that mediate autoimmune encephalomyelitis.** *J Exp Med* 2006, **203**(7):1685-1691.
35. Macatonia SE, Hosken NA, Litton M, Vieira P, Hsieh CS, Culpepper JA, Wysocka M, Trinchieri G, Murphy KM, O'Garra A: **Dendritic cells produce IL-12 and direct the development of Th1 cells from naive CD4+ T cells.** *J Immunol* 1995, **154**(10):5071-5079.
36. Lambrecht BN, Hammad H: **Taking our breath away: dendritic cells in the pathogenesis of asthma.** *Nat Rev Immunol* 2003, **3**(12):994-1003.
37. Brocke R, Riedinger M, Karjalainen K: **Targeted expression of major histocompatibility complex (MHC) class II molecules demonstrates that dendritic cells can induce negative but not positive selection of thymocytes in vivo.** *J Exp Med* 1997, **185**(3):541-550.
38. Jung S, Unutmaz D, Wong P, Sano G, De los Santos K, Sparwasser T, Wu S, Vuthoori S, Ko K, Zavala F, et al: **In vivo depletion of CD11c(+) dendritic cells abrogates priming of CD8(+) T cells by exogenous cell-associated antigens.** *Immunity* 2002, **17**(2):211-220.
39. Barnden MJ, Allison J, Heath WR, Carbone FR: **Defective TCR expression in transgenic mice constructed using cDNA-based alpha- and beta-chain genes under the control of heterologous regulatory elements.** *Immunol Cell Biol* 1998, **76**(1):34-40.
40. Poynter ME, Irvin CG, Janssen-Heininger YM: **Rapid activation of nuclear factor-kappaB in airway epithelium in a murine model of allergic airway inflammation.** *Am J Pathol* 2002, **160**(4):1325-1334.
41. Allard JB, Poynter ME, Marr KA, Cohn L, Rincon M, Whittaker LA: **Aspergillus fumigatus generates an enhanced Th2-biased immune response in mice with defective cystic fibrosis transmembrane conductance regulator.** *J Immunol* 2006, **177**(8):5186-5194.
42. Foster WM, Walters DM, Longphre M, Macri K, Miller LM: **Methodology for the measurement of mucociliary function in the mouse by scintigraphy.** *J Appl Physiol* 2001, **90**(3):1111-1117.
43. Ito T, Wang YH, Duramad O, Hori T, Deslespess GJ, Watanabe N, Qin FX, Yao Z, Cao W, Liu YJ: **TSLP-activated dendritic cells induce an inflammatory T helper type 2 cell response through OX40 ligand.** *J Exp Med* 2005, **202**(9):1213-1223.
44. Jahnsen FL, Moloney ED, Hogan T, Upham JW, Burke CM, Holt PG: **Rapid dendritic cell recruitment to the bronchial mucosa of patients with atopic asthma in response to local allergen challenge.** *Thorax* 2001, **56**(11):823-826.
45. Jahnsen FL, Strickland DH, Thomas JA, Tobagus IT, Napoli S, Zosky GR, Turner DJ, Sly PD, Stumbles PA, Holt PG: **Accelerated antigen sampling and transport by airway mucosal dendritic cells following inhalation of a bacterial stimulus.** *J Immunol* 2006, **177**(9):5861-5867.
46. Richter A, Burrows JP, Nuss H, Granier C, Niemeier U: **Increase in tropospheric nitrogen dioxide over China observed from space.** *Nature* 2005, **437**(7055):129-132.
47. Chen B, Kan H: **Air pollution and population health: a global challenge.** *Environ Health Prev Med* 2008, **13**(2):94-101.
48. Hansbro NG, Horvat JC, Wark PA, Hansbro PM: **Understanding the mechanisms of viral induced asthma: New therapeutic directions.** *Pharmacol Ther* 2008, **117**(3):313-353.
49. Kato M, Hayashi Y, Kimura H: **Oxygen radicals in inflammation and allergy related to viral infections.** *Curr Drug Targets Inflamm Allergy* 2005, **4**(4):497-501.
50. Corry DB, Grunig G, Hadeiba H, Kurup VP, Warnock ML, Sheppard D, Rennick DM, Locksley RM: **Requirements for allergen-induced airway hyperreactivity in T and B cell-deficient mice.** *Mol Med* 1998, **4**(5):344-355.
51. Eisenbarth SC, Piggott DA, Huleatt JW, Visintin I, Herrick CA, Bottomly K: **Lipopolysaccharide-enhanced, toll-like receptor 4-dependent T helper cell type 2 responses to inhaled antigen.** *J Exp Med* 2002, **196**(12):1645-1651.
52. Wilson RH, Whitehead GS, Nakano H, Free ME, Kolls JK, Cook DN: **Allergic sensitization through the airway primes Th17-dependent neutrophilia and airway hyperresponsiveness.** *Am J Respir Crit Care Med* 2009, **180**(8):720-730.
53. Redecke V, Hacker H, Datta SK, Fermin A, Pitha PM, Broide DH, Raz E: **Cutting edge: activation of Toll-like receptor 2 induces a Th2 immune response and promotes experimental asthma.** *J Immunol* 2004, **172**(5):2739-2743.
54. Lambert AL, Dong W, Selgrade MK, Gilmour MI: **Enhanced allergic sensitization by residual oil fly ash particles is mediated by soluble metal constituents.** *Toxicol Appl Pharmacol* 2000, **165**(1):84-93.
55. Lambert AL, Dong W, Winsett DW, Selgrade MK, Gilmour MI: **Residual oil fly ash exposure enhances allergic sensitization to house dust mite.** *Toxicol Appl Pharmacol* 1999, **158**(3):269-277.
56. Diaz-Sanchez D, Garcia MP, Wang M, Jyrala M, Saxon A: **Nasal challenge with diesel exhaust particles can induce sensitization to a neoallergen in the human mucosa.** *J Allergy Clin Immunol* 1999, **104**(6):1183-1188.
57. Whitekus MJ, Li N, Zhang M, Wang M, Horwitz MA, Nelson SK, Horwitz LD, Brechun N, Diaz-Sanchez D, Nel AE: **Thiol antioxidants inhibit the adjuvant effects of aerosolized diesel exhaust particles in a murine model for ovalbumin sensitization.** *J Immunol* 2002, **168**(5):2560-2567.
58. Samuelsen M, Nygaard UC, Lovik M: **Allergy adjuvant effect of particles from wood smoke and road traffic.** *Toxicology* 2008, **246**(2-3):124-131.
59. Trimble NJ, Botelho FM, Bauer CM, Fattouh R, Stampfli MR: **Adjuvant and anti-inflammatory properties of cigarette smoke in murine allergic airway inflammation.** *Am J Respir Cell Mol Biol* 2009, **40**(1):38-46.
60. de Haar C, Hassing I, Bol M, Bleumink R, Pieters R: **Ultrafine carbon black particles cause early airway inflammation and have adjuvant activity in a mouse allergic airway disease model.** *Toxicol Sci* 2005, **87**(2):409-418.
61. Clarke AH, Thomas WR, Rolland JM, Dow C, O'Brien RM: **Murine allergic respiratory responses to the major house dust mite allergen Der p 1.** *Int Arch Allergy Immunol* 1999, **120**(2):126-134.
62. Hammad H, Charbonnier AS, Duez C, Jacquet A, Stewart GA, Tonnel AB, Pestel J: **Th2 polarization by Der p 1-pulsed monocyte-derived dendritic cells is due to the allergic status of the donors.** *Blood* 2001, **98**(4):1135-1141.
63. Wong CK, Li ML, Wang CB, Ip WK, Tian YP, Lam CW: **House dust mite allergen Der p 1 elevates the release of inflammatory cytokines and expression of adhesion molecules in co-culture of human eosinophils and bronchial epithelial cells.** *Int Immunol* 2006, **18**(8):1327-1335.

64. Bleck B, Tse DB, Curotto de Lafaille MA, Zhang F, Reibman J: **Diesel exhaust particle-exposed human bronchial epithelial cells induce dendritic cell maturation and polarization via thymic stromal lymphopoietin.** *J Clin Immunol* 2008, **28**(2):147-156.
65. Bleck B, Tse DB, Jaspers I, Curotto de Lafaille MA, Reibman J: **Diesel exhaust particle-exposed human bronchial epithelial cells induce dendritic cell maturation.** *J Immunol* 2006, **176**(12):7431-7437.
66. Chan RC, Wang M, Li N, Yanagawa Y, Onoe K, Lee JJ, Nel AE: **Pro-oxidative diesel exhaust particle chemicals inhibit LPS-induced dendritic cell responses involved in T-helper differentiation.** *J Allergy Clin Immunol* 2006, **118**(2):455-465.
67. de Haar C, Kool M, Hassing I, Bol M, Lambrecht BN, Pieters R: **Lung dendritic cells are stimulated by ultrafine particles and play a key role in particle adjuvant activity.** *J Allergy Clin Immunol* 2008, **121**(5):1246-1254.
68. Koike E, Kobayashi T: **Ozone exposure enhances antigen-presenting activity of interstitial lung cells in rats.** *Toxicology* 2004, **196**(3):217-227.
69. Porter M, Karp M, Killedear S, Bauer SM, Guo J, Williams D, Breyse P, Georas SN, Williams MA: **Diesel-enriched particulate matter functionally activates human dendritic cells.** *Am J Respir Cell Mol Biol* 2007, **37**(6):706-719.
70. Stumbles PA, Strickland DH, Pimm CL, Proksch SF, Marsh AM, McWilliam AS, Bosco A, Tobagus I, Thomas JA, Napoli S, et al: **Regulation of dendritic cell recruitment into resting and inflamed airway epithelium: use of alternative chemokine receptors as a function of inducing stimulus.** *J Immunol* 2001, **167**(1):228-234.
71. Malavia NK, Raub CB, Mahon SB, Brenner M, Panettieri RA Jr, George SC: **Airway Epithelium Stimulates Smooth Muscle Proliferation.** *Am J Respir Cell Mol Biol* 2009, **41**(3):297-304.
72. Vabulas RM, Ahmad-Nejad P, Ghose S, Kirschning CJ, Issels RD, Wagner H: **HSP70 as endogenous stimulus of the Toll/interleukin-1 receptor signal pathway.** *J Biol Chem* 2002, **277**(17):15107-15112.
73. O'Neill LA, Kaltschmidt C: **NF-kappa B: a crucial transcription factor for glial and neuronal cell function.** *Trends Neurosci* 1997, **20**(6):252-258.
74. Ather JL, Alcorn JF, Brown AL, Guala AS, Suratt BT, Janssen-Heininger YM, Poynter ME: **Distinct Functions of Airway Epithelial NF-[kappa]B Activity Regulate Nitrogen Dioxide-induced Acute Lung Injury.** *Am J Respir Cell Mol Biol* 2009.
75. Ather JL, Hodgkins SR, Janssen-Heininger YM, Poynter ME: **Airway Epithelial NF-[kappa]B Activation Promotes Allergic Sensitization to an Innocuous Inhaled Antigen.** *Am J Respir Cell Mol Biol* 2010.
76. Sung SS, Fu SM, Rose CE Jr, Gaskin F, Ju ST, Beaty SR: **A major lung CD103 (alphaE)-beta7 integrin-positive epithelial dendritic cell population expressing Langerin and tight junction proteins.** *J Immunol* 2006, **176**(4):2161-2172.
77. Kaminsky DA, van der Vliet A, Janssen-Heininger Y: **Reactive nitrogen species in refractory asthma: markers or players?** *J Allergy Clin Immunol* 2008, **121**(2):338-340.
78. Persinger RL, Poynter ME, Ckless K, Janssen-Heininger YM: **Molecular mechanisms of nitrogen dioxide induced epithelial injury in the lung.** *Mol Cell Biochem* 2002, **234-235**(1-2):71-80.
79. Postlethwait EM, Bidani A: **Mechanisms of pulmonary NO<sub>2</sub> absorption.** *Toxicology* 1994, **89**(3):217-237.

doi:10.1186/1465-9921-11-102

**Cite this article as:** Hodgkins et al.: NO<sub>2</sub> inhalation induces maturation of pulmonary CD11c<sup>+</sup> cells that promote antigenspecific CD4<sup>+</sup> T cell polarization. *Respiratory Research* 2010 **11**:102.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

