

RESEARCH

Open Access

# Simvastatin inhibits TGF $\beta$ 1-induced fibronectin in human airway fibroblasts

Dedmer Schaafsma<sup>1,2,3</sup>, Karol D McNeill<sup>1,2</sup>, Mark M Mutawe<sup>1,2</sup>, Saeid Ghavami<sup>1,2,4</sup>, Helmut Unruh<sup>5</sup>, Eric Jacques<sup>6</sup>, Michel Laviolette<sup>6</sup>, Jamila Chakir<sup>6</sup> and Andrew J Halayko<sup>1,2,3,4\*</sup>

## Abstract

**Background:** Bronchial fibroblasts contribute to airway remodelling, including airway wall fibrosis. Transforming growth factor (TGF)- $\beta$ 1 plays a major role in this process. We previously revealed the importance of the mevalonate cascade in the fibrotic response of human airway smooth muscle cells. We now investigate mevalonate cascade-associated signaling in TGF $\beta$ 1-induced fibronectin expression by bronchial fibroblasts from non-asthmatic and asthmatic subjects.

**Methods:** We used simvastatin (1-15  $\mu$ M) to inhibit 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase which converts HMG-CoA to mevalonate. Selective inhibitors of geranylgeranyl transferase-1 (GGT1; GGTI-286, 10  $\mu$ M) and farnesyl transferase (FT; FTI-277, 10  $\mu$ M) were used to determine whether GGT1 and FT contribute to TGF $\beta$ 1-induced fibronectin expression. In addition, we studied the effects of co-incubation with simvastatin and mevalonate (1 mM), geranylgeranylpyrophosphate (30  $\mu$ M) or farnesylpyrophosphate (30  $\mu$ M).

**Results:** Immunoblotting revealed concentration-dependent simvastatin inhibition of TGF $\beta$ 1 (2.5 ng/ml, 48 h)-induced fibronectin. This was prevented by exogenous mevalonate, or isoprenoids (geranylgeranylpyrophosphate or farnesylpyrophosphate). The effects of simvastatin were mimicked by GGTI-286, but not FTI-277, suggesting fundamental involvement of GGT1 in TGF $\beta$ 1-induced signaling. Asthmatic fibroblasts exhibited greater TGF $\beta$ 1-induced fibronectin expression compared to non-asthmatic cells; this enhanced response was effectively reduced by simvastatin.

**Conclusions:** We conclude that TGF $\beta$ 1-induced fibronectin expression in airway fibroblasts relies on activity of GGT1 and availability of isoprenoids. Our results suggest that targeting regulators of isoprenoid-dependent signaling holds promise for treating airway wall fibrosis.

**Keywords:** airway fibroblasts, airway remodeling, asthma, fibronectin, geranylgeranyl transferase, statins

## Background

Chronic obstructive airways diseases, including asthma and COPD, are characterized by structural alterations of the airway wall. The accumulation of extracellular matrix (ECM) proteins (fibrosis) and augmentation of the airway mesenchymal layer, including fibroblasts and airway smooth muscle, are common features of this airway remodeling [1-3]. In asthma, the degree of subepithelial fibrosis has been shown to be associated with disease severity and correlated with a decline in lung

function parameters [4]. Transforming growth factor  $\beta$ 1 (TGF $\beta$ 1) is a principal mediator of subepithelial fibrosis and is highly expressed in asthmatics [4-6]. Airway fibroblasts and myofibroblasts are a primary source of ECM proteins, including fibronectin, in subepithelial fibrosis linked to airway remodeling [7]. Targeting and understanding molecular mechanisms that drive the pro-fibrotic potential of these cells is of great interest with respect to the development of therapies for chronic airways diseases.

Statins were initially developed to inhibit the activity of 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase and are widely prescribed to reduce hyperlipidemia [8]. Substantial evidence shows that statins also

\* Correspondence: ahalayk@cc.umanitoba.ca

<sup>1</sup>Departments of Physiology & Internal Medicine, and Section of Respiratory Disease, University of Manitoba, Winnipeg, MB, Canada

Full list of author information is available at the end of the article

have pleiotropic anti-inflammatory, anti-fibroproliferative and immunomodulatory effects that are independent of their cholesterol-lowering capacity [9-14]. HMG-CoA reductase is the proximal rate-limiting enzyme of the multistep mevalonate cascade for cholesterol biosynthesis. Cholesterol intermediates include the 15- and 20-carbon isoprenoids, farnesylpyrophosphate (FPP) and geranylgeranylpyrophosphate (GGPP), respectively. These lipid moieties are substrates for farnesyl transferase (FT) and geranylgeranyl transferase 1 (GGT1) that catalyze the modification of monomeric G-proteins, such as Ras and RhoA, by conjugating lipid anchors crucial for their association with and activation at the plasma membrane. Effects of statins on cell physiology have been attributed, in part, to the depletion of isoprenoids and the ensuing effects on prenylation-dependent intracellular signaling activity [15-18]. Given the biological importance of FT and GGT1, a number of selective inhibitors have been developed and tested in clinical trials for treatment of cancer [19-21]. To date the impact of these inhibitors on lung health has not been established.

In previous work, we showed that mevalonate-derived isoprenoids provide key regulatory input for the fibrotic

response of human airway smooth muscle cells [14]. We now investigate the role of mevalonate cascade-associated cell signaling in TGF $\beta$ 1-induced expression of the extracellular matrix protein fibronectin by bronchial fibroblasts from both non-asthmatic and asthmatic subjects.


## Materials and methods


### Materials

All chemicals were obtained from Sigma (St. Louis, MO) unless indicated otherwise. Primary antibodies against fibronectin (sc-9068, rabbit polyclonal), collagen type I (sc-8786, goat polyclonal), GGTase 1 $\beta$  (sc-100820, mouse monoclonal) and FT  $\beta$  (sc-137, rabbit polyclonal) were from Santa Cruz Biotechnology (Santa Cruz, CA).

### Human airway fibroblast cell culture: standard study design

Primary human airway fibroblasts were isolated from macroscopically healthy segments of second- to fourth-generation main bronchi obtained after lung resection surgery from patients with a diagnosis of adenocarcinoma. The airway smooth muscle and mesenchymal fibroblast layers were carefully separated by manual dissection; passage 3-4 fibroblasts were used (Figures 1, 2,






and 3). For comparative studies (Figure 4) primary fibroblasts were isolated from bronchial biopsies of mild steroid naïve asthmatic (n = 3) and healthy (n = 3) subjects. The asthmatic subjects fulfilling the American Thoracic Society criteria for asthma [22] were recruited from the Asthma Clinic at IUCPQ (Québec, Canada). They used only an inhaled β2-agonist on demand. The asthmatics were atopic nonsmokers (mean age = 24 ± 2, FEV<sub>1</sub>% predicted = 95 ± 0.4% and PC<sub>20</sub> = 4.6 ± 0.01 mg/ml). None used systemic or inhaled CS. Healthy subjects (mean age = 22 ± 0.4, FEV<sub>1</sub>% predicted = 106 ± 0.82% and PC<sub>20</sub> ≥ 128 mg/ml) were non-atopic nonsmokers with no history of asthma or other pulmonary or systemic diseases. The atopic status of asthmatics was determined by skin prick tests showing a positive reaction (3 mm or more) to at least 2 aero-allergens. The healthy group had no skin reaction. Bronchial biopsies were obtained by bronchoscopy from asthmatic and healthy subjects as described previously [23]; passage 4-6 cells were used. Written informed consent was obtained from all subjects before entry into the study. All procedures were approved by the Human Research Ethics Board (University of Manitoba) and the Ethics

Committee at the Institut Universitaire de Cardiologie et de Pneumologie de Québec.

Cells were plated on uncoated plastic dishes in Dulbecco's modified Eagle's medium (DMEM) supplemented with 50 U/ml streptomycin, 50 μg/ml penicillin, and 10% fetal bovine serum (FBS). Cells were grown to ~80% confluence, after which they were maintained for 24 hours in serum-free DMEM supplemented with 5 μg/ml insulin, 5 μg/ml transferrin, and 5 ng/ml selenium.

For all studies, unless otherwise stated, we followed a standard treatment protocol. Serum-deprived cells were stimulated with TGFβ1 (2.5 ng/ml) for 48 hrs in the presence or absence of simvastatin (1, 10, 15 μM, added 30 min before TGFβ1). In some experiments, the effects of co-incubation with mevalonic acid (1 mM) [14,24], geranylgeranyl pyrophosphate (GGPP, 30 μM) [14,18] or farnesyl pyrophosphate (FPP, 30 μM) [14,18] were studied (all added 4 h prior to TGFβ1). In separate experiments the effects of the geranylgeranyltransferase inhibitor GGTI-286 (10 μM) [14,18,24,25] and the farnesyltransferase inhibitor FTI-277 (10 μM) [14,25,26] were investigated (added 30 min prior to TGFβ1).




### Protein immunoblotting

After washing cultures with ice-cold phosphate-buffered saline (PBS, composition (mM): NaCl 140.0; KCl 2.6; KH<sub>2</sub>PO<sub>4</sub> 1.4; Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O 8.1; pH 7.4) cell lysates were prepared in ice-cold SDS buffer (composition: 62.5 mM Tris-HCl, 2% SDS, 1 mM PMSF, 1 mM protease inhibitor mix, and 1 mM phosphatase inhibitor mix). Equal amounts of protein, as determined using a commercial Lowry assay, were subjected to electrophoresis and transferred to nitrocellulose membranes. Membranes were subsequently blocked in Tris buffer containing 0.1% Tween-20 and 5% w/v dried milk powder, then incubated overnight at 4°C with primary antibodies (fibronectin (diluted 1:1000), GGTase 1β (diluted 1:400), FTβ (diluted 1:1000) and β-actin (diluted 1:20.000)). Blots were then incubated with diluted horseradish

peroxidase conjugated secondary antibodies prior to visualizing bands on film using enhanced chemiluminescence reagents (Amersham, Buckinghamshire, UK). All blots were subjected to densitometry using a computer page scanner and Totalab™ software. For data analyses bands were normalized to β-actin to correct for small differences in loading.

### RNA extraction and reverse transcriptase PCR

Total RNA was extracted using the RNeasy RNA Mini Kit (Qiagen, U.S.A). For reverse transcription (first strand cDNA synthesis) we used 2 μg of total RNA (in × μL), 0.3 μL Random Hexamers (3 mg/mL, Invitrogen) and 10-× μL ddH<sub>2</sub>O. After heating for 5 min at 65°C, 9 μL of reaction mixture (1 μL dNTP PCR mix (10 mM, Amersham, Canada), 4 μL 5 × first-strand buffer, 2 μL DTT (0.1 M), 1



**Figure 4 Simvastatin profoundly suppresses the augmented TGFβ1-induced fibronectin expression in asthmatic bronchial fibroblasts.**

Western analysis was performed using whole cell lysates obtained from nonasthmatic and asthmatic bronchial fibroblast cultures that were maintained under TGFβ1 (2.5 ng/ml)-stimulated or serum-deprived conditions in the absence or presence of simvastatin (0.1-10 μM) for 48 h. (A) Western blots showing simvastatin dose-dependently suppresses TGFβ1-induced fibronectin expression in nonasthmatic and asthmatic bronchial fibroblasts, whereas GGase 1β and FTase β protein abundance is not affected. (B) Representative western blot showing a more robust TGFβ1-induced fibronectin abundance in asthmatic versus nonasthmatic fibroblasts, which is effectively suppressed by simvastatin. (C) Densitometric analysis revealed significant differences in TGFβ1-induced fibronectin expression between nonasthmatic and asthmatic fibroblasts. Moreover, simvastatin markedly prevents fibronectin expression in both cell types. Data represent means ± s.e. mean of 5 independent experiments, using cells obtained from 3 non-asthmatic and 3 asthmatic subjects. *Legend*: t = 0: initial basal (unstimulated) cultures; t = 48 h: treatment time-matched, basal (unstimulated) cultures; Simv: simvastatin. \*\*P < 0.01, \*\*\*P < 0.001 compared to asthmatic TGFβ. #P < 0.05, ###P < 0.001 compared to nonasthmatic TGFβ.



μL RNaseOUT (40 U) and 1 μL Moloney murine leukemia virus reverse transcriptase (M-MLV RT, 200 U, Invitrogen) was added. Samples were incubated at 42°C for 120 minutes then heating at 72°C for 15 minutes. cDNA was stored at -20°C until further use. PCR amplification was performed in a total volume of 50 μL which included 1 μL RT reaction mixture, 0.5 μM of each forward and reverse oligonucleotide, 1 × PCR buffer with 1.5 mM MgCl, 0.2 mM dNTP PCR mix and 1.25 U of Platinum Taq Polymerase (Invitrogen). Primers used for GAPDH and the human prenyltransferase subunits FNTA (var1), FNTA (var2), FNTB, PGGT1B, RabGGTA and RabGGTB are listed in Table 1.

### Statistical analysis

All data represent means ± s.e. mean from n separate experiments. Statistical significance of differences was evaluated by the Student's *t*-test for paired observations or by ANOVA for multiple measurements followed by a Tukey's post-test. Differences were considered to be statistically significant when *P* < 0.05.

## Results

### Simvastatin prevents TGFβ1-induced fibronectin protein expression

Primary human bronchial mesenchymal fibroblasts were stimulated with 2.5 ng/ml TGFβ1 for 48 h in the presence and absence of simvastatin (Figures 1A and 1B). TGFβ1 induced a marked increase in fibronectin protein, an effect significantly suppressed by 1 (69.5 ± 7.4% of C), 10 (62.5 ± 6.7% of C) and 15 μM (37.6 ± 11.5% of C) simvastatin. Similarly, TGFβ1-induced collagen I protein abundance was dose-dependently inhibited by simvastatin (Figure 1A), indicating that as for airway smooth muscle [14] the inhibitory effects of simvastatin are more broadly applicable. Based on these data and previous reports by our group on potential toxicity of high concentrations of simvastatin [27], we used 10 μM in all subsequent experiments.

### Depletion of isoprenoids underpins the suppressive effects of simvastatin

To determine whether the effects of simvastatin on fibronectin are due to reduced formation of mevalonate,

FPP and GGPP, we incubated human airway fibroblasts with TGFβ1 and simvastatin in the presence of mevalonate (1 mM), FPP (30 μM) or GGPP (30 μM). Co-incubation with these intermediates caused nearly full prevention of the suppressive effects of simvastatin, implying their depletion is critical for the effects of simvastatin (Figures 2A and 2B).

### Inhibition of GGT1, but not FT, mimics the effects of simvastatin

We next investigated the effects of the geranylgeranyl transferase inhibitor GGTI-286 (10 μM) and the farnesyl transferase inhibitor FTI-277 (10 μM) on TGFβ1-induced fibronectin protein expression (Figures 3A and 3B). GGTI-286 significantly prevented TGFβ1-induced fibronectin accumulation to a similar degree as 10 μM simvastatin. In contrast, no reduction in fibronectin was observed after co-treatment with FTI-277 (Figures 3A and 3B). These findings indicate a predominant involvement of GGT1, but not FT, in the TGFβ1-induced profibrotic response of human airway fibroblasts. In line with these findings, profiling of the expression of protein prenyltransferase subunits by RT-PCR revealed expression of 6 subunits, including two variants of the farnesyltransferase, CAAX box, alpha (FNTA) subunit that is common to both GGT1 and FT (Figure 3C). These results indicate human airway fibroblasts express the genes necessary to form GGT1, FT and GGT2 prenyltransferase heterodimers. Further confirming these findings, we demonstrate that GGTase 1β and FTase β protein are expressed in non-asthmatic and asthmatic fibroblasts; abundance of these subunits was not affected by simvastatin, nor was there any difference in expression between non-asthmatic and asthmatic fibroblasts (Figure 4A).

### Simvastatin effectively suppresses the augmented profibrotic response of asthmatic bronchial fibroblasts

To determine the effects of simvastatin on fibronectin expression in non-asthmatic and asthmatic bronchial fibroblasts, cells were stimulated with TGFβ1 in the presence and absence of simvastatin (Figure 4A). Simvastatin dose-dependently suppressed fibronectin abundance in non-asthmatic and asthmatic fibroblasts. To directly

**Table 1 Primers Used For Reverse Transcriptase PCR**

Fragment	Forward	Reverse	PCR Product Size
FNTA var 1(Human)	5'-TAT AGA TCC GGT GCC GCA GAA TGA-3'	5'-ACT CTC CGG AAA TGC CAC ACT GTA-3'	196 bp
FNTA var 2 (Human)	5'-GTC CTG CAG CGT GAT GAA AGA AGT-3'	5'-ACT CTC CGG AAA TGC CAC ACT GTA-3'	101 bp
FNTB (Human)	5'-TGC AGA GGG AGA AGC ACT TCC ATT-3'	5'-AGC TGT GCA GGA TCC AAT AGC AGA-3'	114 bp
PGGT1B (Human)	5'-TTG CAA TGA CCT ACA CTG GCC TCT-3'	5'-TCA CTG CCT TCA GGT ACT GCA CAA-3'	143 bp
RabGGTA (Human)	5'-TGC TGG AGA ATA GCG TGC TCA AGA-3'	5'-AGT CAA GAT GGG TGA CCA AGA GCA-3'	121 bp
RabGGTB (Human)	5'-AGA CCA GGT TCT GAA TCC CAT GCT-3'	5'-TGG TAA CTT CTC CGG CCT TCC ATT-3'	162 bp
GAPDH (Human)	5'-AGC AAT GCC TCC TGC ACC ACC AAC-3'	5'-CCG GAG GGG CCA TCC ACA GTC T-3'	136 bp

compare these responses different lysates were analyzed on the same gel; though no differences in basal fibronectin expression were observed, a more robust response to TGF $\beta$ 1 by asthmatic fibroblasts was evident ( $2.7 \pm 0.4$  fold higher; Figures 4B and 4C). Importantly, simvastatin suppressed TGF $\beta$ 1-induced fibronectin expression in both non-asthmatic and asthmatic cells ( $40.0 \pm 4.8\%$  and  $52.4 \pm 8.1\%$  reduction, respectively; Figure 4C).

## Discussion

In the present study, we demonstrate that isoprenoid intermediates of the mevalonate cascade provide key regulatory input for the TGF $\beta$ 1-induced expression of the extracellular matrix protein fibronectin by human bronchial fibroblasts. HMG-CoA reductase inhibition with simvastatin suppressed TGF $\beta$ 1-induced fibronectin abundance, an effect prevented by exogenous mevalonate, GGPP and FPP. Effects of simvastatin were mirrored by the selective GGT1 inhibitor, GGTI-286, but not the farnesyl protein transferase inhibitor, FTI-277, suggesting that proteins targeted by GGT1 for conjugation of prenyl lipid chains are crucial for TGF $\beta$ 1-induced fibronectin expression. Moreover, we show for the first time that fibronectin expression in response to TGF $\beta$ 1 is markedly augmented in bronchial fibroblasts obtained from asthmatics compared to those from non-asthmatics. Simvastatin effectively inhibited TGF $\beta$ 1-induced fibronectin in fibroblasts from both groups.

Statins are recognized for pleiotropic effects that exceed their cholesterol lowering capacity. Statin use correlates with reduced COPD hospitalizations and mortality [28-30], and up to 50% slower decline in lung function (FEV1 and FVC) in smokers, former smokers and non-smokers [9,10]. In patients receiving double lung transplant, statin use is associated with significantly better post-operative spirometry and airway inflammation as indicated by reduced numbers of neutrophils and lymphocytes [31]. Several recent studies have also revealed anti-inflammatory effects of statins in murine and rat models of allergic asthma [32,33] and COPD [11,12]. Moreover, statins reportedly suppress *ex vivo* airway responsiveness in animal models [34,35].

Statins have broad effects on cell responses, including inhibition of proliferation, migration and they can promote apoptosis [15-18,27]. These studies are consistent with our observation that mevalonate, GGPP and FPP can prevent the effects of simvastatin, confirming the fundamental role of regulated protein lipidation in cell function, including fibronectin expression [36]. Importantly, we have demonstrated previously that under the conditions studied 10  $\mu$ M simvastatin does not affect human airway fibroblast viability, as determined by

MTT assays, within 48 h; indicating the observed decrease in fibronectin is not an artifact due to cell death [27]. Our finding that mevalonate, FPP and GGPP prevent the suppressive effects of simvastatin yet only GGTI-286, but not FTI-277, mimics its actions suggests that signaling proteins that are subject to GGT1-catalyzed geranylgeranylation are crucial for TGF $\beta$ 1-induced fibronectin expression in airway fibroblasts. These findings are supported by studies using human fetal lung fibroblasts demonstrating the effectiveness of a GGT1 inhibitor (GGTI-298), but not a FT inhibitor (FTI-277), on TGF $\beta$ 1-mediated expression of connective tissue growth factor (CTGF), elastin and fibronectin mRNA [21,37-39].

The lack of effect of FT inhibition versus the effectiveness of FPP to prevent the inhibitory effects of simvastatin seems paradoxical. Theoretically, FPP can be converted to GGPP intracellularly, as such providing a substrate for GGT1 [40]. Although an interesting hypothesis, in the presence of simvastatin, even with the addition of FPP, formation of the more downstream sterol intermediate GGPP is not effected as HMG-CoA inhibition depletes the upstream 5-carbon upstream intermediate, isopentyl pyrophosphate (IPP), that is required for conversion of FPP to GGPP [40]. An alternative potential explanation could lie in the promiscuity of GGT1 both in using alternate isoprenoids (i.e. FPP) and in effectively prenylating downstream effectors that are essential for TGF $\beta$ 1-induced fibronectin expression. Thus, our findings suggest that GGT1 may be able to utilize FPP to modify a critical downstream effector. Moreover, we speculate that FT is unable to prenylate signaling proteins and induce their activation when GGT1 activity is suppressed with GGTI-286. These complex topics need to be addressed mechanistically in future studies.

The anti-fibrotic effects of statins are not likely to be limited to airway mesenchymal cells. Indeed, beneficial effects of statins on human hypertrophic cardiomyopathy [41] and the occurrence of renal interstitial fibrosis in transgenic rabbits [42] have been reported. In addition, statins have cardioprotective effects that are associated with their anti-fibrotic effects in adrenomedullin-knockout mice [43] and have been reported to prevent left ventricular remodelling, including interstitial fibrosis, in hypertensive rats [44]. *In vitro* studies using human lung fibroblasts derived from healthy and idiopathic pulmonary fibrosis (IPF) patients also demonstrate that simvastatin can inhibit connective tissue growth factor expression, reduce collagen gel contraction, and down-regulate smooth muscle  $\alpha$ -actin expression [45]. In addition, systemic administration of simvastatin markedly attenuates the onset of collagen-associated lung fibrosis in mice treated with trachea-instilled bleomycin [46].

To our knowledge, we demonstrate for the first time that TGF $\beta$ 1-induced fibronectin protein expression is significantly greater in fibroblasts from asthmatic subjects compared to those obtained from healthy subjects. These results correlate well with findings by Westergren-Thorsson and colleagues that demonstrate fibroblasts isolated from asthmatics produce increased amounts of proteoglycans [47]. This intrinsic difference between asthmatic and non-asthmatic fibroblasts to express ECM proteins could contribute to sub-epithelial fibrosis in the asthmatic airway. Our data indicate that fibronectin expression by asthmatic fibroblasts is not-refractory to simvastatin, suggesting this therapeutic approach could be of benefit. In clinical studies, short-term treatment of asthmatics with statins had no significant effect on lung function or other indices of asthma control in patients treated with corticosteroids [48] or without anti-inflammatory medication [49]. Conversely, a recent study revealed that simvastatin can enhance the anti-inflammatory effects of inhaled corticosteroids in mild asthmatics [50], which is in line with reduced alveolar macrophage numbers in sputum of asthmatics that had received statin treatment [48]. Inasmuch as these studies indicate that the effects of short-term statin treatment on airway inflammation and lung function in mild to moderate asthmatics is debatable, the effects of statins on features of airway remodelling, which are generally associated with disease duration and severity, remain elusive. Recent in vitro studies using human airway smooth muscle cells and fibroblasts do show statins inhibit proliferation and promote apoptosis [18,51], which when considered in the context of previous work by our group [14] and the present study showing a concomitant effect on fibronectin expression in bronchial mesenchymal cells, suggests potential for suppressing airway remodeling.

## Conclusions

Our data indicate that mevalonate cascade associated cell signaling is a key signaling component in TGF $\beta$ 1-induced fibronectin expression in primary human airway fibroblasts. Moreover, it appears that the prenyltransferase GGT1 is a principal effector for isoprenoid-dependent TGF $\beta$ 1 induced fibronectin expression. Last, we demonstrate the presence of exaggerated fibronectin expression in response to TGF $\beta$ 1 in asthmatic fibroblasts, and confirm that simvastatin can significantly suppress the response in these cells. Based on our results simvastatin and perhaps more selective inhibitors of GGT1 could be considered as potential therapeutic tools to modulate airway wall fibrosis in fibrotic airway diseases such as asthma.

## Abbreviations

ECM: extracellular matrix; FPP: farnesylpyrophosphate; FT: farnesyl transferase; GGPP: geranylgeranyl pyrophosphate; GGTase: geranylgeranyl transferase;

HMG-CoA: 3-hydroxy-3-methylglutaryl-coenzyme A; TGF: transforming growth factor.

## Acknowledgements

This work was supported by grants from the Canadian Institutes of Health Research (CIHR), GlaxoSmithKline Collaborative Innovation Research Fund, Manitoba Institute of Child Health, and Canada Foundation for Innovation. DS holds a Canadian Institutes of Health Research (CIHR) Postdoctoral fellowship, and is supported by CIHR/HSFC IMPACT Grant in Pulmonary and Cardiovascular Research. SG holds a Parker B. Francis postdoctoral fellowship. AJH holds a Canada Research Chair in Airway Cell and Molecular Biology.

## Author details

<sup>1</sup>Departments of Physiology & Internal Medicine, and Section of Respiratory Disease, University of Manitoba, Winnipeg, MB, Canada. <sup>2</sup>Biology of Breathing Theme, Manitoba Institute of Child Health, Winnipeg, MB, Canada. <sup>3</sup>CIHR IMPACT Training Program in Pulmonary and Cardiovascular Research, University of Manitoba, Winnipeg, MB, Canada. <sup>4</sup>CIHR National Training Program in Allergy and Asthma, University of Manitoba, Winnipeg, MB, Canada. <sup>5</sup>Section of Thoracic Surgery, University of Manitoba, Winnipeg, MB, Canada. <sup>6</sup>Institut Universitaire de Cardiologie et de Pneumologie de Québec, Québec, Canada.

## Authors' contributions

DS participated in the design and coordination of the study, performed a major part of the experiments, performed the statistical analysis and drafted the manuscript. KDM, MMM and SG substantially assisted in performing the experiments. EJ, HU, and ML were responsible for tissue collection and handling. JC participated in interpretation of results and the preparation of the manuscript. AJH supervised the study, participated in its design and data interpretation, and was involved in the manuscript preparation. All authors read and approved the final manuscript.

## Competing interests

The authors declare that they have no competing interests.

Received: 8 June 2011 Accepted: 24 August 2011

Published: 24 August 2011

## References

1. An SS, Bai TR, Bates JH, Black JL, Brown RH, Brusasco V, Chitano P, Deng L, Dowell M, Eidelman DH, et al: **Airway smooth muscle dynamics: a common pathway of airway obstruction in asthma.** *Eur Respir J* 2007, **29**:834-860.
2. Jeffery PK: **Remodeling in asthma and chronic obstructive lung disease.** *Am J Respir Crit Care Med* 2001, **164**:S28-38.
3. Postma DS, Timens W: **Remodeling in asthma and chronic obstructive pulmonary disease.** *Proc Am Thorac Soc* 2006, **3**:434-439.
4. Minshall EM, Leung DY, Martin RJ, Song YL, Cameron L, Ernst P, Hamid Q: **Eosinophil-associated TGF-beta1 mRNA expression and airways fibrosis in bronchial asthma.** *Am J Respir Cell Mol Biol* 1997, **17**:326-333.
5. Holgate ST, Holloway J, Wilson S, Howarth PH, Haitchi HM, Babu S, Davies DE: **Understanding the pathophysiology of severe asthma to generate new therapeutic opportunities.** *J Allergy Clin Immunol* 2006, **117**:496-506, quiz 507.
6. Redington AE, Roche WR, Madden J, Frew AJ, Djukanovic R, Holgate ST, Howarth PH: **Basic fibroblast growth factor in asthma: measurement in bronchoalveolar lavage fluid basally and following allergen challenge.** *J Allergy Clin Immunol* 2001, **107**:384-387.
7. Matsumoto H, Niimi A, Takemura M, Ueda T, Minakuchi M, Tabuena R, Chin K, Mio T, Ito Y, Muro S, et al: **Relationship of airway wall thickening to an imbalance between matrix metalloproteinase-9 and its inhibitor in asthma.** *Thorax* 2005, **60**:277-281.
8. Nakamura H, Arakawa K, Itakura H, Kitabatake A, Goto Y, Toyota T, Nakaya N, Nishimoto S, Muranaka M, Yamamoto A, et al: **Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial.** *Lancet* 2006, **368**:1155-1163.
9. Alexeeff SE, Litonjua AA, Sparrow D, Vokonas PS, Schwartz J: **Statin use reduces decline in lung function: VA Normative Aging Study.** *Am J Respir Crit Care Med* 2007, **176**:742-747.



10. Keddissi JI, Younis WG, Chbeir EA, Daher NN, Dernaika TA, Kinasewitz GT: **The use of statins and lung function in current and former smokers.** *Chest* 2007, **132**:1764-1771.
11. Lee JH, Lee DS, Kim EK, Choe KH, Oh YM, Shim TS, Kim SE, Lee YS, Lee SD: **Simvastatin inhibits cigarette smoking-induced emphysema and pulmonary hypertension in rat lungs.** *Am J Respir Crit Care Med* 2005, **172**:987-993.
12. Lee SD, Lee JH, Kim EK, Choi KH, Oh YM, Shim TS: **Effects of simvastatin on cigarette smoking-induced structural and functional changes in rat lungs.** *Chest* 2005, **128**:574S.
13. Liao JK, Laufs U: **Pleiotropic effects of statins.** *Annu Rev Pharmacol Toxicol* 2005, **45**:89-118.
14. Schaafsma D, Dueck G, Ghavami S, Kroeker A, Mutawe MM, Hauff K, Xu FY, McNeill KD, Unruh H, Hatch GM, Halayko AJ: **The Mevalonate Cascade as a Target to Suppress Extracellular Matrix Synthesis by Human Airway Smooth Muscle.** *Am J Respir Cell Mol Biol* 2011, **44**:394-403.
15. Guijarro C, Blanco-Colio LM, Ortego M, Alonso C, Ortiz A, Plaza JJ, Diaz C, Hernandez G, Egido J: **3-Hydroxy-3-methylglutaryl coenzyme a reductase and isoprenylation inhibitors induce apoptosis of vascular smooth muscle cells in culture.** *Circ Res* 1998, **83**:490-500.
16. Kim DY, Ryu SY, Lim JE, Lee YS, Ro JY: **Anti-inflammatory mechanism of simvastatin in mouse allergic asthma model.** *Eur J Pharmacol* 2007, **557**:76-86.
17. Liao JK: **Isoprenoids as mediators of the biological effects of statins.** *J Clin Invest* 2002, **110**:285-288.
18. Takeda N, Kondo M, Ito S, Ito Y, Shimokata K, Kume H: **Role of RhoA inactivation in reduced cell proliferation of human airway smooth muscle by simvastatin.** *Am J Respir Cell Mol Biol* 2006, **35**:722-729.
19. Eskens FA, Awada A, Cutler DL, de Jonge MJ, Luyten GP, Faber MN, Statkevich P, Sparreboom A, Verweij J, Hanauske AR, Piccart M: **Phase I and pharmacokinetic study of the oral farnesyl transferase inhibitor SCH 66336 given twice daily to patients with advanced solid tumors.** *J Clin Oncol* 2001, **19**:1167-1175.
20. Sebt SM, Hamilton AD: **Farnesyltransferase and geranylgeranyltransferase I inhibitors and cancer therapy: lessons from mechanism and bench-to bedside translational studies.** *Oncogene* 2000, **19**:6584-6593.
21. Zujewski J, Horak ID, Bol CJ, Woestenborghs R, Bowden C, End DW, Piotrovsky VK, Chiao J, Belly RT, Todd A, et al: **Phase I and pharmacokinetic study of farnesyl protein transferase inhibitor R115777 in advanced cancer.** *J Clin Oncol* 2000, **18**:927-941.
22. **Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease (COPD) and asthma. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, November 1986.** *Am Rev Respir Dis* 1987, **136**:225-244.
23. Goulet F, Boulet LP, Chakir J, Tremblay N, Dube J, Laviolette M, Boutet M, Xu W, Germain L, Auger FA: **Morphologic and functional properties of bronchial cells isolated from normal and asthmatic subjects.** *Am J Respir Cell Mol Biol* 1996, **15**:312-318.
24. Fujimoto M, Oka T, Murata T, Hori M, Ozaki H: **Fluvastatin inhibits mast cell degranulation without changing the cytoplasmic Ca<sup>2+</sup> level.** *Eur J Pharmacol* 2009, **602**:432-438.
25. Lesh RE, Emala CW, Lee HT, Zhu D, Panettieri RA, Hirshman CA: **Inhibition of geranylgeranylation blocks agonist-induced actin reorganization in human airway smooth muscle cells.** *Am J Physiol Lung Cell Mol Physiol* 2001, **281**:L824-831.
26. Nonaka M, Uota S, Saitoh Y, Takahashi M, Sugimoto H, Amet T, Arai A, Miura O, Yamamoto N, Yamaoka S: **Role for protein geranylgeranylation in adult T-cell leukemia cell survival.** *Exp Cell Res* 2009, **315**:141-150.
27. Ghavami S, Mutawe MM, Hauff K, Stelmack GL, Schaafsma D, Sharma P, McNeill KD, Hynes TS, Kung SK, Unruh H, et al: **Statin-triggered cell death in primary human lung mesenchymal cells involves p53-PUMA and release of Smac and Omi but not cytochrome c.** *Biochim Biophys Acta* 2010, **1803**:452-467.
28. Frost FJ, Petersen H, Tollestrup K, Skipper B: **Influenza and COPD mortality protection as pleiotropic, dose-dependent effects of statins.** *Chest* 2007, **131**:1006-1012.
29. Mancini GB, Etminan M, Zhang B, Levesque LE, FitzGerald JM, Brophy JM: **Reduction of morbidity and mortality by statins, angiotensin-converting enzyme inhibitors, and angiotensin receptor blockers in patients with chronic obstructive pulmonary disease.** *J Am Coll Cardiol* 2006, **47**:2554-2560.
30. Soyseth V, Brekke PH, Smith P, Omland T: **Statin use is associated with reduced mortality in COPD.** *Eur Respir J* 2007, **29**:279-283.
31. Johnson BA, Iacono AT, Zeevi A, McCurry KR, Duncan SR: **Statin use is associated with improved function and survival of lung allografts.** *Am J Respir Crit Care Med* 2003, **167**:1271-1278.
32. McKay A, Leung BP, McInnes IB, Thomson NC, Liew FY: **A novel anti-inflammatory role of simvastatin in a murine model of allergic asthma.** *J Immunol* 2004, **172**:2903-2908.
33. Zeki AA, Franzi L, Last J, Kenyon NJ: **Simvastatin inhibits airway hyperreactivity: implications for the mevalonate pathway and beyond.** *Am J Respir Crit Care Med* 2009, **180**:731-740.
34. Chiba Y, Arima J, Sakai H, Misawa M: **Lovastatin inhibits bronchial hyperresponsiveness by reducing RhoA signaling in rat allergic asthma.** *Am J Physiol Lung Cell Mol Physiol* 2008, **294**:L705-713.
35. Chiba Y, Sato S, Misawa M: **Inhibition of antigen-induced bronchial smooth muscle hyperresponsiveness by lovastatin in mice.** *J Smooth Muscle Res* 2008, **44**:123-128.
36. Morck C, Olsen L, Kurth C, Persson A, Storm NJ, Svensson E, Jansson JO, Hellqvist M, Enejder A, Faergeman NJ, Pilon M: **Statins inhibit protein lipidation and induce the unfolded protein response in the non-sterial producing nematode *Caenorhabditis elegans*.** *Proc Natl Acad Sci USA* 2009, **106**:18285-18290.
37. Kucich U, Rosenbloom JC, Herrick DJ, Abrams WR, Hamilton AD, Sebt SM, Rosenbloom J: **Signaling events required for transforming growth factor-beta stimulation of connective tissue growth factor expression by cultured human lung fibroblasts.** *Arch Biochem Biophys* 2001, **395**:103-112.
38. Kucich U, Rosenbloom JC, Shen G, Abrams WR, Blaskovich MA, Hamilton AD, Ohkanda J, Sebt SM, Rosenbloom J: **Requirement for geranylgeranyl transferase I and acyl transferase in the TGF-beta-stimulated pathway leading to elastin mRNA stabilization.** *Biochem Biophys Res Commun* 1998, **252**:111-116.
39. Kucich U, Rosenbloom JC, Shen G, Abrams WR, Hamilton AD, Sebt SM, Rosenbloom J: **TGF-beta1 stimulation of fibronectin transcription in cultured human lung fibroblasts requires active geranylgeranyl transferase I, phosphatidylcholine-specific phospholipase C, protein kinase C-delta, and p38, but not erk1/erk2.** *Arch Biochem Biophys* 2000, **374**:313-324.
40. Swanson KM, Hohl RJ: **Anti-cancer therapy: targeting the mevalonate pathway.** *Curr Cancer Drug Targets* 2006, **6**:15-37.
41. Patel R, Nagueh SF, Tsybouleva N, Abdellatif M, Lutucuta S, Kopelen HA, Quinones MA, Zoghbi WA, Entman ML, Roberts R, Marian AJ: **Simvastatin induces regression of cardiac hypertrophy and fibrosis and improves cardiac function in a transgenic rabbit model of human hypertrophic cardiomyopathy.** *Circulation* 2001, **104**:317-324.
42. Vieira JM Jr, Mantovani E, Rodrigues LT, Delle H, Noronha IL, Fujihara CK, Zatz R: **Simvastatin attenuates renal inflammation, tubular transdifferentiation and interstitial fibrosis in rats with unilateral ureteral obstruction.** *Nephrol Dial Transplant* 2005, **20**:1582-1591.
43. Yamamoto C, Fukuda N, Jumabay M, Saito K, Matsumoto T, Ueno T, Soma M, Matsumoto K, Shimosawa T: **Protective effects of statin on cardiac fibrosis and apoptosis in adrenomedullin-knockout mice treated with angiotensin II and high salt loading.** *Hypertens Res* 2010.
44. Zhao XY, Li L, Zhang JY, Liu GQ, Chen YL, Yang PL, Liu RY: **Atorvastatin prevents left ventricular remodeling in spontaneously hypertensive rats.** *Int Heart J* 2010, **51**:426-431.
45. Watts KL, Sampson EM, Schultz GS, Spiteri MA: **Simvastatin inhibits growth factor expression and modulates profibrogenic markers in lung fibroblasts.** *Am J Respir Cell Mol Biol* 2005, **32**:290-300.
46. Ou XM, Feng YL, Wen FQ, Huang XY, Xiao J, Wang K, Wang T: **Simvastatin attenuates bleomycin-induced pulmonary fibrosis in mice.** *Chin Med J (Engl)* 2008, **121**:1821-1829.
47. Westergren-Thorsson G, Chakir J, Lafreniere-Allard MJ, Boulet LP, Tremblay GM: **Correlation between airway responsiveness and proteoglycan production by bronchial fibroblasts from normal and asthmatic subjects.** *Int J Biochem Cell Biol* 2002, **34**:1256-1267.
48. Hotherhall EJ, Chaudhuri R, McSharry C, Donnelly I, Lafferty J, McMahon AD, Weir CJ, Meiklejohn J, Sattar N, McInnes I, et al: **Effects of atorvastatin added to inhaled corticosteroids on lung function and sputum cell counts in atopic asthma.** *Thorax* 2008, **63**:1070-1075.

49. Menzies D, Nair A, Meldrum KT, Fleming D, Barnes M, Lipworth BJ: **Simvastatin does not exhibit therapeutic anti-inflammatory effects in asthma.** *J Allergy Clin Immunol* 2007, **119**:328-335.
50. Maneechotesuwan K, Ekjitrakul W, Kasetsinsombat K, Wongkajornsilp A, Barnes PJ: **Statins enhance the anti-inflammatory effects of inhaled corticosteroids in asthmatic patients through increased induction of indoleamine 2, 3-dioxygenase.** *J Allergy Clin Immunol* 2010, **126**:754-762 e751.
51. Ghavami S, Mutawe MM, Hauff K, Stelmack GL, Schaafsma D, Sharma P, McNeill KD, Hynes TS, Kung SK, Unruh H, *et al*: **Statin-triggered cell death in primary human lung mesenchymal cells involves p53-PUMA and release of Smac and Omi but not cytochrome c.** *Biochim Biophys Acta* 2010.

doi:10.1186/1465-9921-12-113

**Cite this article as:** Schaafsma *et al.*: Simvastatin inhibits TGF $\beta$ 1-induced fibronectin in human airway fibroblasts. *Respiratory Research* 2011 **12**:113.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

